

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 March 2001 (22.03.2001)

PCT

(10) International Publication Number
WO 01/19866 A1

- (51) International Patent Classification⁷: C07K 14/47, C12N 9/64, 15/12, C12Q 1/37, A61K 38/48, A61P 19/02, 31/18, 37/00, 37/06, G01N 33/53
- (21) International Application Number: PCT/AU00/01085
- (22) International Filing Date:
11 September 2000 (11.09.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
PQ 2762 10 September 1999 (10.09.1999) AU
PQ 5709 18 February 2000 (18.02.2000) AU
- (71) Applicant (for all designated States except US): THE UNIVERSITY OF SYDNEY [AU/AU]; Parramatta Road, Sydney, NSW 2006 (AU).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): ABBOTT, Catherine, Anne [AU/AU]; 288 Trafalgar Street, Annandale, NSW 2038 (AU). GORELL, Mark, Douglas [AU/AU]; 9 Piper Street North, Annandale, NSW 2038 (AU).
- (74) Agent: GRIFFITH HACK; GPO Box 4164, Sydney, NSW 2001 (AU).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— With international search report.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/19866 A1

(54) Title: DIPEPTIDYL PEPTIDASES

(57) Abstract: A novel human dipeptidyl aminopeptidase (DPP8) with structural and functional similarities to DPPIV and fibroblast activation protein (FAP) has been identified and characterized. The novel peptide comprises the sequence ID NO: 1, the amino acid sequences His⁷³⁶GlyTrpSerTyrGlyGlyTyrLeu, Leu⁸¹⁶AspGluAsnValHisPheAlaHis, Glu⁸⁴⁷ArgHisSerIleArg, Phe²²⁵ValLeuGlnGluGluPhe, or a sequence having at least 60 % homology with, and the substrate specificity of SEQ ID NO:1. A gene encoding the dipeptidyl peptidase and uses of the dipeptidyl peptidase.

- 1 -

TITLE

DIPEPTIDYL PEPTIDASES

FIELD OF INVENTION

- 5 The invention relates to a dipeptidyl peptidase, to a nucleic acid molecule which encodes it, and to uses of the peptidase.

BACKGROUND OF THE INVENTION

- 10 The dipeptidyl peptidase (DPP) IV-like gene family is a family of molecules which have related protein structure and function [1-3]. The gene family includes the following molecules: DPPIV (CD26), dipeptidyl amino-peptidase-like protein (DPP6) and fibroblast activation
15 protein (FAP) [1,2,4,5]. Another possible member is DPPIV- β [6].

The molecules of the DPPIV-like gene family are serine proteases, they are members of the peptidase family S9b,
20 and together with prolyl endopeptidase (S9a) and acylaminoacyl peptidase (S9c), they are comprised in the prolyl oligopeptidase family [5,7].

DPPIV and FAP both have similar postproline dipeptidyl
25 amino peptidase activity, however, unlike DPPIV, FAP also has gelatinase activity [8,9].

DPPIV substrates include chemokines such as RANTES, eotaxin, macrophage-derived chemokine and stromal-cell-
30 derived factor 1; growth factors such as glucagon and glucagon-like peptides 1 and 2; neuropeptides including neuropeptide Y and substance P; and vasoactive peptides [10-12].

35 DPPIV and FAP also have non-catalytic activity; DPPIV binds adenosine deaminase, and FAP binds to $\alpha_3\beta_1$ and $\alpha_5\beta_1$ integrin [13-14].

- 2 -

In view of the above activities, the DPPIV-like family members are likely to have roles in intestinal and renal handling of proline containing peptides, cell adhesion, peptide metabolism, including metabolism of cytokines, 5 neuropeptides, growth factors and chemokines, and immunological processes, specifically T cell stimulation[3,11,12].

Consequently, the DPPIV-like family members are likely to 10 be involved in the pathology of disease, including for example, tumour growth and biology, type II diabetes, cirrhosis, autoimmunity, graft rejection and HIV infection[3,15-18].

15 Inhibitors of DPPIV have been shown to suppress arthritis, and to prolong cardiac allograft survival in animal models *in vivo*[19,20]. Some DPPIV inhibitors are reported to inhibit HIV infection[21]. It is anticipated that DPPIV inhibitors will be useful in other therapeutic 20 applications including treating diarrhoea, growth hormone deficiency, lowering glucose levels in non insulin dependent diabetes mellitus and other disorders involving glucose intolerance, enhancing mucosal regeneration and as immunosuppressants[3,21-24].

25 There is a need to identify members of the DPPIV-like gene family as this will allow the identification of inhibitor(s) with specificity for particular family member(s), which can then be administered for the purpose 30 of treatment of disease. Alternatively, the identified member may of itself be useful for the treatment of disease.

35

- 3 -

SUMMARY OF THE INVENTION

The present invention seeks to address the above identified need and in a first aspect provides a peptide which comprises the amino acid sequence shown in SEQ ID NO:1.

This peptide has substrate specificity for the following compounds: H-Ala-Pro-pNA, H-Gly-Pro-pNA and H-Arg-Pro-pNA. Therefore, it is a prolyl oligopeptidase and a dipeptidyl
10 peptidase, because it is capable of hydrolysing the peptide bond C-terminal to proline in each of these compounds.

The peptide is homologous with human DPPIV, and
15 importantly, identity between the sequences of DPPIV and SEQ ID NO: 1 is observed at the region of DPPIV containing the catalytic triad residues and the two glutamate residues of the β -propeller domain essential for DPPIV enzyme activity. The observation of amino acid sequence
20 homology means that the peptide which has the amino acid sequence shown in SEQ ID NO:1 is a member of the DPPIV-like gene family. Accordingly the peptide was provisionally named DPPIVL1, and is now named and described herein as DPP8.

25

The following sequences of the human DPPIV amino acid sequence are important for the catalytic activity of DPPIV: (i) Tyr⁶²⁷GlyTrpSerTyrGlyGlyTyrVal; (ii) Ala⁷⁰⁷AspAspAsnValHisPhe; (iii) Glu⁷³⁸AspHisGlyIleAlaGln;
30 and (iv) Tyr²⁰¹ValTyrGluGluGluVal [25-28]. As described herein, the alignment of the following sequences of DPP8: His⁷³⁶GlyTrpSerTyrGlyGlyTyrLeu; Leu⁸¹⁶AspGluAsnValHisPheAla; Glu⁸⁴⁷ArgHisSerIleArg and Phe²⁵⁵ValLeuGlnGluGluPhe with
35 these sequences of DPP8 are likely to confer the catalytic activity of DPP8. Thus, in a second aspect, the invention provides a peptide comprising the following amino acid

- 4 -

sequences: His⁷³⁶GlyTrpSerTyrGlyGlyTyrLeu;
Leu⁸¹⁶AspGluAsnValHisPheAlaHis; Glu⁸⁴⁷ArgHisSerIleArg and
Phe²⁵⁵ValLeuGlnGluGluPhe; which has the substrate
specificity of the sequence shown in SEQ ID NO:1.

5

Also described herein, using multiple sequence alignment,
it is observed that DPP8 has 55% amino acid similarity and
32% amino acid identity with a *C. elegans* protein.

Further, as shown herein, a nucleic acid molecule which
10 encodes DPP8, is capable of hybridising specifically with
DPP8 sequences derived from non-human species. Together
these data suggest that DPP8 is expressed in non-human
species. Thus in a third aspect, the invention provides a
peptide which has at least 60% amino acid identity with
15 the amino acid sequence shown in SEQ ID NO:1, and which
has the substrate specificity of the sequence shown in SEQ
ID NO:1. Preferably, the amino acid identity is 75%.

More preferably, the amino acid identity is 95%. Amino
acid identity is calculated using GAP software [GCC
20 Version 8, Genetics Computer Group, Madison, WI, USA] as
described further herein. Typically, the non-human DPP8
comprises the following sequences:

His⁷³⁶GlyTrpSerTyrGlyGlyTyrLeu;
Leu⁸¹⁶AspGluAsnValHisPheAlaHis; Glu⁸⁴⁷ArgHisSerIleArg and
25 Phe²⁵⁵ValLeuGlnGluGluPhe.

In view of the homology between DPPIV and DPP8 amino acid
sequences, it is expected that these sequences will have
similar tertiary structure. This means that the tertiary
30 structure of DPP8 is likely to include the seven-blade β -
propeller domain and the α/β hydrolase domain of DPPIV.
These structures in DPP8 are likely to be conferred by the
regions comprising β -propeller, Gly¹⁸⁰ to Asp⁶⁰⁶, α/β
hydrolase, Ser⁶⁰⁷ to Ile⁸⁸² and about 70 to 100 residues in
35 the region Arg³⁹ to Gln¹⁷⁹. As it is known that the β -
propeller domain regulates proteolysis mediated by the
catalytic triad in the α/β hydrolase domain of prolyl

- 5 -

oligopeptidase, [29] it is expected that truncated forms of DPP8 can be produced, which have the substrate specificity of the sequence shown in SEQ ID NO:1, comprising the regions referred to above

5 (His⁷³⁶GlyTrpSerTyrGlyGlyTyrLeu; Leu⁸¹⁶AspGluAsnValHisPheAlaHis; Glu⁸⁴⁷ArgHisSerIleArg and Phe²⁵⁵ValLeuGlnGluGluPhe) which confer the catalytic specificity of DPP8. Examples of truncated forms of DPP8 which might be prepared are those in which the region

10 conferring the β -propeller domain and the α/β hydrolase domain are spliced together. Other examples of truncated forms include those which are encoded by splice variants of DPP8 mRNA. Thus although, as described herein, the biochemical characterisation of DPP8 shows that DPP8

15 consists of 882 amino acids and has a molecular weight of about 100kDa, it is recognised that truncated forms of DPP8 which have the substrate specificity of the sequence shown in SEQ ID NO:1, may be prepared using standard techniques [30,31]. Thus in a fourth aspect, the

20 invention provides a fragment of the sequence shown in SEQ ID NO: 1, which has the substrate specificity of the sequence shown in SEQ ID NO:1. Preferably, the fragment has an amino acid sequence shown in SEQ ID NO: 3, 5 or 7.

25 As described herein, the sequence shown in SEQ ID NO:1 does not contain a consensus sequence for N-linked glycosylation. Therefore it is unlikely that DPP8 is associated with N-linked glycosylation. In this regard, DPP8 is distinguished from other DPPIV-like gene family

30 members, which contain between 6 and 9 consensus sequences for N-linked glycosylation. Thus in one embodiment, an asparagine residue in the peptide of the first aspect of the invention is not linked to a carbohydrate molecule.

35 The analysis of DPP8 expression described herein shows that it is likely that DPP8 is expressed as a cytoplasmic protein. The expression of DPP8 is therefore

- 6 -

distinguished from other DPPIV-like gene family members, which are expressed on the cytoplasmic membrane, or in other words, the cell surface membrane. Thus in another embodiment, the peptide of the first aspect of the invention is not expressed on a cell surface membrane of a cell.

It is recognised that DPP8 may be fused, or in other words, linked to a further amino acid sequence, to form a fusion protein which has the substrate specificity of the sequence shown in SEQ ID NO:1. An example of a fusion protein is described herein which comprises the sequence shown in SEQ ID NO:1 which is linked to a further amino acid sequence: a "tag" sequence which consists of an amino acid sequence encoding the V5 epitope and a His tag. An example of another further amino acid sequence which may be linked with DPP8 is a glutathione S transferase (GST) domain [30]. Another example of a further amino acid sequence is a portion of CD8 α [8]. Thus in one aspect, the invention provides a fusion protein comprising the amino acid sequence shown in SEQ ID NO:1 linked with a further amino acid sequence, the fusion protein having the substrate specificity of the sequence shown in SEQ ID NO:1.

25

It is also recognised that the peptide of the first aspect of the invention may be comprised in a polypeptide, so that the polypeptide has the substrate specificity of DPP8. The polypeptide may be useful, for example, for altering the protease susceptibility of DPP8, when used in *in vivo* applications. An example of a polypeptide which may be useful in this regard, is albumin. Thus in another embodiment, the peptide of the first aspect is comprised in a polypeptide which has the substrate specificity of DPP8.

35

- 7 -

As described above, the isolation and characterisation of DPP8 is necessary for identifying inhibitors of DPP8 catalytic activity, which may be useful for the treatment of disease. A method for identifying inhibitors of DPP8 catalytic activity, described herein, has identified that various inhibitors of DPPIV and serine proteases, zinc and mimetic peptides, Ala-Pro-Gly and Lys-Pro, but not inhibitors of metalloproteinases, aspartyl proteinases or cysteinyl proteinases, inhibit DPP8 catalytic activity. Accordingly, in a fifth aspect, the invention provides a method of identifying a molecule capable of inhibiting cleavage of a substrate by DPP8, the method comprising the following steps:

- (a) contacting DPP8 with the molecule;
- (b) contacting DPP8 of step (a) with a substrate capable of being cleaved by DPP8, in conditions sufficient for cleavage of the substrate by DPP8; and
- (c) detecting substrate not cleaved by DPP8, to identify that the molecule is capable of inhibiting cleavage of the substrate by DPP8.

It is recognised that although inhibitors of DPP8 may also inhibit DPPIV and other serine proteases, as described herein, the alignment of the DPP8 amino acid sequence with most closely related molecules, (i.e. DPPIV), reveals that the DPP8 amino acid is distinctive, particularly at the regions controlling substrate specificity. Accordingly, it is expected that it will be possible to identify inhibitors which inhibit DPP8 catalytic activity specifically, which do not inhibit catalytic activity of DPPIV-like gene family members, or other serine proteases. Thus, in a sixth aspect, the invention provides a method of identifying a molecule capable of inhibiting specifically, the cleavage of a substrate by DPP8, the method comprising the following steps:

- 8 -

(a) contacting DPP8 and a further protease with the molecule;

(b) contacting DPP8 and the further protease of step (a) with a substrate capable of being cleaved by DPP8 and the further protease, in conditions sufficient for cleavage of the substrate by DPP8 and the further protease; and

(c) detecting substrate not cleaved by DPP8, but cleaved by the further protease, to identify that the molecule is capable of inhibiting specifically, the cleavage of the substrate by DPP8.

In a seventh aspect, the invention provides a method of reducing or inhibiting the catalytic activity of DPP8, the method comprising the step of contacting DPP8 with an inhibitor of DPP8 catalytic activity. As various inhibitors of DPPIV catalytic activity are shown herein to inhibit DPP8 catalytic activity, it is recognised that other inhibitors of DPPIV may be useful for inhibiting DPP8 catalytic activity. Examples of inhibitors suitable for use in the seventh aspect are described in [21,32,33]. Other inhibitors useful for inhibiting DPP8 catalytic activity can be identified by the methods of the fifth or sixth aspects of the invention, which methods are exemplified herein.

In one embodiment, the catalytic activity of DPP8 is reduced or inhibited in a mammal by administering the inhibitor of DPP8 catalytic activity to the mammal. It is recognised that these inhibitors have been used to reduce or inhibit DPPIV catalytic activity *in vivo*, and therefore, may also be used for inhibiting DPP8 catalytic activity *in vivo*. Examples of inhibitors useful for this purpose are disclosed in the following [21,32-34].

Preferably, the catalytic activity of DPP8 in a mammal is reduced or inhibited in the mammal, for the purpose of

- 9 -

treating a disease in the mammal. Diseases which are likely to be treated by an inhibitor of DPP8 catalytic activity are those in which DPPIV-like gene family members are associated [3,10,11,17,21,36], including for example, 5 neoplasia, type II diabetes, cirrhosis, autoimmunity, graft rejection and HIV infection.

Preferably, the inhibitor for use in the seventh aspect of the invention is one which inhibits the cleavage of a 10 peptide bond C-terminal adjacent to proline. As described herein, examples of these inhibitors are 4-(2-aminoethyl)benzenesulfonylfluoride, aprotinin, benzamidine/HCl, Ala-Pro-Gly, H-Lys-Pro-OH HCl salt and zinc ions, for example, zinc sulfate or zinc chloride. 15 More preferably, the inhibitor is one which specifically inhibits DPP8 catalytic activity, and which does not inhibit the catalytic activity of other serine proteases, including, for example DPPIV or FAP.

20 In an eighth aspect, the invention provides a method of cleaving a substrate which comprises contacting the substrate with DPP8 in conditions sufficient for cleavage of the substrate by DPP8, to cleave the substrate. Examples of molecules which can be cleaved by the method 25 are H-Ala-Pro-pNA, H-Gly-Pro-pNA and H-Arg-Pro-pNA. The conditions sufficient for cleaving the substrate are described herein. Molecules which are cleaved by DPPIV including RANTES, eotaxin, macrophage-derived chemokine, stromal-cell-derived factor 1, glucagon and glucagon-like 30 peptides 1 and 2, neuropeptide Y, substance P and vasoactive peptide are also likely to be cleaved by DPP8 [11,12]. In one embodiment, the substrate is cleaved by cleaving a peptide bond C-terminal adjacent to proline in the substrate. The molecules cleaved by DPP8 may have 35 Ala, or Trp, Ser, Gly, Val or Leu in the P1 position, in place of Pro [11,12].

- 10 -

As described herein, DPP8 gene expression is upregulated in stimulated lymphocyte and lymphocytic cell lines which suggests that DPP8 may have a functional role in T cell costimulation and proliferation. It is recognised
5 therefore that measuring DPP8 gene expression is useful for detecting T cell activation. Thus in a ninth aspect, the invention provides a method of detecting an activated T cell, the method comprising the step of detecting the level of DPP8 gene expression in a T cell. In one
10 embodiment, the level of DPP8 gene expression is detected by measuring the amount of DPP8 mRNA in the cell, as described herein.

The inventors have characterised the sequence of a nucleic
15 acid molecule which encodes the amino acid sequence shown in SEQ ID NO:1. Thus in a tenth aspect, the invention provides a nucleic acid molecule which encodes the amino acid sequence shown in SEQ ID NO:1.

20 In an eleventh aspect, the invention provides a nucleic acid molecule which consists of the sequence shown in SEQ ID NO:2.

As described herein, at least three splice variants of
25 DPP8 RNA which have an open reading frame from 2.6 to 3.1 kb in length are observed. As a frame shift mutation or termination signal was not observed in the sequence of these splice variants, and as the coding sequence of two of the splice variants includes a sequence which encodes
30 the amino acid sequence associated with catalytic activity, it is recognised that some of the peptides encoded by the splice variants are likely to have the substrate specificity of DPP8. Thus in an embodiment, the nucleic acid molecule is a fragment of the sequence shown
35 in SEQ ID NO: 1 which is about 2.6 to 3.1 kb in length and which encodes a peptide which has the substrate

- 11 -

specificity of the sequence shown in SEQ ID NO:1.

Preferably, the nucleic acid molecule has a sequence shown in any one of SEQ ID NO.s: 4, 6 and 8.

5 In a twelfth aspect, the invention provides a nucleic acid molecule which is capable of hybridising to a nucleic acid molecule consisting of the sequence shown in SEQ ID NO:2 in stringent conditions, and which encodes a peptide which
10 ID NO:1. As shown in the Northern blot analysis described herein, DPP8 mRNA hybridises specifically to the sequence shown in SEQ ID NO:2, after washing in 2XSSC/ 1.0%SDS at 37°C, or after washing in 0.1XSSC/0.1% SDS at 50°C.

"Stringent conditions" are conditions in which the nucleic
15 acid molecule is exposed to 2XSSC/ 1.0% SDS. Preferably, the nucleic acid molecule is capable of hybridising to a molecule consisting of the sequence shown in SEQ ID NO:2 in high stringent conditions. "High stringent conditions" are conditions in which the nucleic acid molecule is
20 exposed to 0.1XSSC/ 0.1%SDS at 50°C.

As described herein, the inventors believe that the gene which encodes DPP8 is located at band q22 on human chromosome 15. The location of the DPP8 gene is
25 distinguished from genes encoding other prolyl oligopeptidases, which are located on chromosome 2, at bands 2q24.3 and 2q23, or chromosome 7. Thus in an embodiment, the nucleic acid molecule is one capable of hybridising to a gene which is located at band q22 on
30 human chromosome 15.

It is recognised that a nucleic acid molecule which encodes the amino acid sequence shown in SEQ ID NO:1, or which comprises has the sequence shown in SEQ ID NO:2,
35 could be made by producing the fragment of the sequence which is translated, using standard techniques [30,31].

- 12 -

Thus in an embodiment, the nucleic acid molecule does not contain 5' or 3' untranslated sequences.

In a thirteenth aspect, the invention provides a vector
5 which comprises a nucleic acid molecule of the tenth
aspect of the invention. In one embodiment, the vector is
capable of replication in a COS-7 cell, CHO cell or 293T
cell, or E.coli. In another embodiment, the vector is
selected from the group consisting of λ TripleEx, pTripleEx,
10 pGEM-T Easy Vector, pSecTag2Hygro, pet15b, pEE14.HCMV.gs
and pCDNA3.1/V5/His.

In a fourteenth aspect, the invention provides a cell
which comprises a vector of the thirteenth aspect of the
15 invention. In one embodiment, the cell is an E.coli cell.
Preferably, the E. coli is MC1061, DH5 α , JM109, BL21DE3,
pLyss. In another embodiment, the cell is a COS-7, COS-1,
293T or CHO cell.

20 In a fifteenth aspect, the invention provides a method for
making a peptide of the first aspect of the invention
comprising, maintaining a cell according to the fourteenth
aspect of the invention in conditions sufficient for
expression of the peptide by the cell. The conditions
25 sufficient for expression are described herein. In one
embodiment, the method comprises the further step of
isolating the peptide.

In a sixteenth aspect, the invention provides a peptide
30 when produced by the method of the fifteenth aspect.

In a seventeenth aspect, the invention provides a
composition comprising a peptide of the first aspect and a
pharmaceutically acceptable carrier.

35

- 13 -

In an eighteenth aspect, the invention provides an antibody which is capable of binding a peptide according to the first aspect of the invention. The antibody can be prepared by immunising a subject with purified DPP8 or a
5 fragment thereof according to standard techniques [35]. As described herein, an antibody was prepared by immunising with transiently transfected DPP8⁺ cells. It is recognised that the antibody is useful for inhibiting activity of DPP8, or for detecting increased gene
10 expression of DPP8, for the purpose of identifying an activated T cell. In one embodiment, the antibody of the eighth aspect of the invention is produced by a hybridoma cell.

15 In a nineteenth aspect, the invention provides a hybridoma cell which secretes an antibody of the nineteenth aspect.

BRIEF DESCRIPTION OF THE FIGURES

20 Figure 1. Cloning strategy for isolating full-length DPP8 cDNA and the alternative splicing variants of DPP8 observed. Representation of three splice variants is shown including loss of serine recognition site by one splice variant (T8).

25 Figure 2. Nucleotide sequence and amino acid sequence of human DPP8. The nucleotide and predicted one letter code amino acid sequence are shown. This sequence shows no putative membrane spanning domain (deduced from
30 hydrophobicity plots) or potential N-linked glycosylation sites. The putative serine recognition site and aspartic acid and histidine which form the Ser-Asp-His catalytic triad are marked. Base pairs are numbered in the right margin.

35

- 14 -

Figure 3. Alignment of the deduced amino acid residue sequence of DPP8 with the *C. elegans* homolog of DPP8 and human DPPIV. Amino-acid residues are numbered in the right margin. Amino-acid residues identical in all three proteins are boxed. Asterisks mark the putative catalytic triad residues and two glutamates of the β -propeller domain essential for DPPIV enzyme activity. The grey shading denotes the α/β hydrolase domain of these proteins. Filled triangles joined by lines indicate starts and ends of alternatively spliced transcripts, stPBMCDy3-3-10 (solid lines), T8 (dashed lines) and T21 (solid lines). The alignment was constructed using the PILEUP program in GCG.

Figure 4. Northern Blot analysis of DPP8 expression. Human multiple tissue Northern blots (CLONTECH) containing 2 μ g per lane of poly A⁺ RNA were hybridized with a ³²P labeled DPP8 probe at 68°C and washed at high stringency. The autoradiograph was exposed for 1 day at -70°C with a BIOMAX MS screen. Molecular mass markers are indicated in base pairs on the left side of each autoradiogram. Figure 4a. Master RNA (CLONTECH) blot of poly A⁺ RNA was hybridized with a ³²P labelled DPP8 probe at 65°C and washed at high stringency. The autoradiograph was exposed for 3 days at -70°C with BIOMAX MS screen. DPP8 mRNA was detected in all tissues examined.

Figure 5. Chromosomal localization of human DPP8. Metaphase showing FISH with the biotinylated DPP8 cDNA probe. Normal male chromosomes stained with DAPI. Hybridization sites on chromosome 15 are indicated by an arrow.

Figure 6. Western blot analysis of transfected cell lines. Analysis of lysates of stable cell lines. DPP8 protein was

- 15 -

seen in DPP8 /V5/His stable cell lines but not in DPP4 or vector-only stable cell lines. The electrophoretic mobility of the protein was not altered when samples were boiled. The band of greater mobility was probably a breakdown product of intact DPP8.

Figure 7. DPP8 enzyme activity. (A) pH-dependence of DPP8 enzyme activity. (B) DPP8 and DPPIV enzyme kinetics. Means +/- SD of absorbance change per minute, multiplied by 1000 are shown. Curve fitting assumed Michaelis-Menten kinetics.

Figure 8. RT-PCR analysis of DPP8 expression. PCR amplifications with primers specific for either a portion of human DPP8 that contained no alternate splicing, Val416 to Gly 679 (top of each gel) or glyceraldehyde-3-phosphate dehydrogenase (G3PDH) (bottom of each gel. (A) Top gel, lanes 1-5 contain PCR products from unstimulated PBMC cDNA from five subjects. Bottom gel, lanes 6 to 11 contain PCR products from OKT3-stimulated PBMC cDNA from six subjects. (B). PCR products are from cDNA from lymphocytic cell lines, liver or placenta as indicated. Negative control amplifications contained reaction mix, enzyme and no cDNA template. Each PCR was performed for 35 cycles. The PCR products were electrophoresed on agarose gels and stained with ethidium bromide. The left lane of each gel contains PUC19 digested with *Hae*III as size markers.

Figure 9. Northern blot analysis of murine DPP8 expression. A murine Northern blot containing 10 µg per lane of total RNA was hybridized with a ³²P-labeled human DPP8 probe at 60°C and washed at low stringency. Autoradiographic exposure was for 3 days at -70°C with a BIOMAX MS screen.

35

- 16 -

DETAILED DESCRIPTION OF THE INVENTION

EXAMPLES

General

- 5 Restriction enzymes and other enzymes used in cloning were obtained from Boehringer Mannheim Roche. Standard molecular biology techniques were used [31] unless indicated otherwise.
- 10 An EST clone (GENBANKTM accession number AA417787) was obtained from American Type Culture Collection. The DNA insert of this clone was sequenced on both strands using automated sequencing at SUPAMAC (Sydney, Australia).

15 Cell culture and RNA preparation

- Human peripheral blood monocytes (PBMCs) were isolated by Ficoll-Hypaque density-gradient centrifugation (Pharmacia, Uppsala, Sweden) of blood obtained from healthy donors. The PBMCs were incubated in AIM-V medium (Life Technologies, Gaithersburg, MD, USA) supplemented with 2 mM L-glutamine and were stimulated with either 1 $\mu\text{g} \cdot \text{mL}^{-1}$ phytohaemagglutinin (Wellcome) or 100 ng $\cdot \text{mL}^{-1}$ OKT3 (Orthoclone, FL, USA) for 72 h. The human cell lines Jurkat, CCRF-CEM, Raji, Daudi and HepG2 were grown to
- 20 confluence in Dulbecco's modified Eagle's medium (Trace Biosciences, NSW, Australia) supplemented with 10% fetal bovine serum and 2 mM L-glutamine.

- Liver and placental RNA were prepared from snap-frozen
- 30 human tissue as described previously [37]. However, RNA was prepared from PBMCs and cell lines using an RNeasy kit (Qiagen, Germany).

Bioinformatics

- 35 BLAST programs [38] and all multiple sequence alignments

- 17 -

were performed through the Australian National Genomic Information Service (ANGIS, Sydney, NSW, Australia). PILEUP (GCG Version 8, Genetics Computer Group, Madison, WI, USA) was used for multiple sequence alignments of
5 proteins.

A BLAST search was performed on the public expressed sequence tag (EST) database using the complete human DPPIV (GenBank™ accession number X60708) and FAP (accession
10 number U09278) nucleotide sequences as query sequences. An EST clone (accession number AA417787) was obtained from the American Type Culture Collection. The DNA insert of this clone was sequenced on both strands using automated sequencing at SUPAMAC (Sydney, NSW, Australia). Because
15 of its homology with DPPIV, this new gene was named dipeptidyl peptidase 8 (DPP8).

DPP8 Cloning

ESTAA417787 was used to design forward (caa ata gaa att
20 gac gat cag gtg) and reverse (tct tga agg tag tgc aaa aga tgc) DPP8 primers for polymerase chain reaction (PCR) from ESTAA417787. The PCR conditions were as follows: 94°C for 5 min, followed by 35 cycles of 94°C for 1 minute, 55°C for 30 sec and 70°C for 1 min. This 484 bp PCR product was gel
25 purified, ³²P-α labelled using Megaprime Labeling Kit (Amersham Pharmacia Biotec, UK) and hybridized to a Master RNA blot (CLONTECH, Palo Alto, CA, USA) that contained poly A⁺ from 50 adult and fetal tissues immobilized in dots as per manufacturers' instructions. This Master RNA blot
30 was also probed with DPP4 for comparison of mRNA tissue expression.

The forward and reverse DPP8 primers were used for PCR to screen a human placental λ STRETCH PLUS library (CLONTECH,
35 Palo Alto, CA, USA) for the presence of DPP8 cDNA in the

- 18 -

library. The library was then screened by standard molecular biology techniques [30,31]. After primary screening, 23 clones were selected for secondary screening, after which 22 remained positive. For the tertiary screen the clones contained in λ TriplEx were converted into pTriplEx plasmids and transformed into BM25.8 *E. coli* recipient bacteria. The plated bacteria were screened and it was confirmed that all 22 clones were positive. Two of these clones, T8 and T21 were selected for further study.

5' RACE (Rapid amplification of cDNA ends)

A 5' RACE Version 2.0 kit (Gibco BRL, Life technologies) was applied on activated T cell (ATC) and placental RNA as prescribed in the kit instructions. The T8 DNA sequence was used to design GSP 1 (TCC TTC CTT CAG CAT CAA TC) and GSP2 (CTT AAA AGT GAC TTT AGG ATT TGC TGT ACC). 5' RACE PCR products were cloned into pGEM-T Easy® Vector (Promega Co., Madison, WI, USA) and sequenced by primer walking.

Confirmation of identity of RACE product

Reverse transcriptase PCR was carried out on ATC RNA using DPP8-pr23 (GGA AGA AGA TGC CAG ATC AGC TGG) and DPP8-pr19r (TCC GTG TAT CCT GTA TCA TAG AAG) to span across the junction between the RACE product and the EST and library clones. Two gel purified products ATCd3-2-1 (1603bp) and ATC3-3-10 (1077bp) were cloned into pGEM-T Easy® (Promega Co., Madison, WI, USA) and sequenced.

Subcloning of DPP8 cDNA into a pcDNA3.1/V5/His Expression Vector

The ATC RACE product, the ATCd3-2-1 (1603bp) junction fragment and the library clone T21 were joined together and cloned into the expression vector pcDNA3.1/V5/His A (Invitrogen, the Netherlands) to form a DPP8 cDNA of 3.1

- 19 -

kb with an open reading frame of 882 aa. The first construct was made using three sequential cloning steps. Firstly, a *Eco RV/Xba I* fragment of T21 (containing 3' DPP8, stop codon and 3' untranslated region on DPP8 cDNA) was ligated into the vector pcDNA3.1/V5/His A which had been digested with *Eco RV/Xba I*. An *Eco RI/Eco RV* fragment of ATCd3-2-1 was then added to this construct digested with *Eco RI/Eco RV*. Finally the RACE product was cut with *Eco RI* and cloned into the *Eco RI* site of the previous construct to form the complete 3.1 kb DPP8 cDNA. This construct pcDNA3.1-DPP8 expressed protein with no detectable tag. In addition the stop codon in the DPP8 expression construct in pcDNA3.1/V5/His V5 was genetically altered using PCR to create a C-terminal fusion with the V5 and His tag contained in the vector. This construct was named pcDNA3.1- DPP8/V5/His. All expression constructs subcloned into pcDNA3.1/V5/His were verified by full sequence analysis.

20 DPP8 gene expression by Northern Blot

Human multiple tissue Northern blots (CLONTECH) containing 2 ug of poly A⁺ RNA were prehybridized in Express Hybridization solution (CLONTECH) for 30 min at 68°C. Both the DPP8 484 bp product and the 5' RACE ATC product were radiolabeled using a Megaprime Labeling kit (Amersham Pharmacia Biotech) and [³²P]dCTP (NEN Dupont). Unincorporated label was removed using a NICK column (Amersham Pharmacia Biotech) and the denatured probe was incubated for 2 hrs at 68°C in Express Hybridization solution. Washes were performed at high stringency and blots exposed to BIOMAX MS film for overnight with a BIOMAX MS screen at -70°C.

DPP8 gene expression in mice by Northern Blot

- 20 -

A Northern blot containing 10 ug of total liver RNA per lane was made using standard methods [31]. The RNA was derived from male and female mice of two strains, C57Bl6 and Balb/c. The Northern blot was prehybridized in Express Hybridization solution (CLONTECH, Palo Alto, USA) for 1 hr at 60°C. A 2.4kb human DPP8 cDNA (PCR product) was radiolabeled using the Megaprime Labeling kit (Amersham Pharmacia Biotech) and [³²P]dCTP (NEN Dupont). Unincorporated label was removed using a NICK column (Amersham Pharmacia Biotech) and the denatured probe was incubated with the blot overnight at 60°C in Express Hybridization solution. Washes were performed at low stringency (2 x SSC/0.05% SDS for 1 h at 37°C followed by 0.1x SSC/0.1% SDS for 30 min at 40°C) and blots exposed to BIOMAX MS film for three days with a BIOMAX MS screen at -70°C.

Expression of DPP8 in mouse liver using rtPCR

Mouse liver RNA was reverse transcribed using the Superscript II enzyme kit (Gibco BRL, Gaithersburg, MD) as described previously [42]. The cDNA was diluted 1 in 4 and stored in aliquots at -70°C. PCR using mouseDPP8-pr1F (atg att acc acc cag gaa gcg) as the forward primer and mouseDPP8-pr2R (atc tcc gac atc ttg aaa gtg acc) as the reverse primer was used to detect mouse DPP8 mRNA.

One ul of diluted cDNA was amplified in a 50 ul PCR reaction which contained: 0.2 mM dNTPs, 1 ul of 50 x Advantage 2 Polymerase Mix (Clontech), 1 X Advantage 2 PCR buffer (Clontech) and 100 ng of each primer. The PCR involved an initial step of 95°C for 1 min to inactivate the TaqStart Antibody. This was followed by 35 cycles; denaturation at 95°C for 30 sec, 68°C for 1 min, followed by a final step of 68°C for 1 min. The amplified products were analysed by electrophoresis of 10 µl of PCR reaction on a 3:1 Nusieve gel (FMC Bioproducts, Rockville, MD) plus

- 21 -

0.5 µg/ml. ethidium bromide in TAE buffer (0.04M Tris acetate, 0.001 M EDTA, pH 8.0). The gel was then Southern Blotted using standard techniques [31]. The Southern blot was hybridized at 60°C for 2hr with the 2.4 kb human DPP8
5 cDNA probe prepared as described above. Washes were performed at low stringency (2 x SSC/0.05% SDS for 1 h at 37°C followed by 0.1x SSC/0.1% SDS for 40 min at 50°C). The blot was exposed to XAR5 Kodak film for 30 min at RT.

10 DPP8 expression by RT-PCR

Reverse transcriptase PCR was performed on human ATC RNA, human placental RNA and human liver RNA using TED primers DPP8/pr3 (GCA CTA CCT TCA AGA AAA CCT TGG) and DPP8/pr20R (TAT GGT ATT GCT GGG TCT CTC AGG) to give a 293 bp
15 product.

Transfection, Western blot, immunocytochemistry, cytochemistry and flow cytometry

Monkey kidney fibroblast (COS-7) cells (American Type
20 Culture Collection, CRL-1651) were grown and transfected as described previously [39]. For making stable cell lines, Geneticin (G418; Gibco-BRL) was added to the medium, beginning 24 h after transfection. COS cell extracts were prepared by sonication followed by
25 differential centrifugation and neither boiled nor reduced before SDS/PAGE (10% gel) and transfer to nitrocellulose, as described previously [40,9]. The presence of DPP8 fused with the V5 epitope was detected using an anti-V5 mAb (Invitrogen). COS cell monolayers were fixed in cold
30 ethanol before staining with anti-V5 mAb [39,41,9]. Some monolayers were fixed in 4% paraformaldehyde and permeabilized with 0.1% Triton X-100 [35], then double-stained with wheat germ agglutinin to label Golgi apparatus and with goat anti-mouse IgG to label DPP8,
35 conjugated to Alexa Fluor 488 and Alexa Fluor 594,

- 22 -

respectively (Molecular Probes, Eugene, OR, USA). Flow cytometry and confocal scanning microscopy using a Leica TCS-NT confocal microscope have been described previously [39,9].

5

Purification of recombinant DPP8/V5/His and DPPIV/V5/His
Cells (1×10^7) expressing each protein were sonicated in native buffer (50 mM sodium phosphate, 300 mM NaCl), then
10 treated with 700 U DNase for 20 min at room temperature. DPPIV is expressed at the cell surface, so 1% Triton X-100 was used to solubilize DPPIV/V5/His. Insoluble material was removed by centrifugation. The supernatant was incubated with 1 mL Talon® Metal Affinity Resin (Clontech)
15 following the manufacturer's instructions for a batch/gravity flow procedure. The resin was washed with 50 mM sodium phosphate, containing 300 mM NaCl and 5 mM imidazole, and proteins were eluted using the same buffer containing 150 mM imidazole. Enzyme activity was used to
20 monitor eluted fractions.

Enzyme assays

Enzyme assays were performed as described previously [1]. Either clarified cell extract from 1×10^4 sonicated COS-7
25 cells or purified protein derived from 1×10^5 cells was incubated with substrate in 70 μ L phosphate buffer, pH 7.4, for 30 min at 37°C, except where otherwise indicated. The specific DPPIV substrates, Gly-Pro-toluenesulfonate, H-Gly-Pro-p-nitroanilide (NA)/HCl (Sigma, St Louis, MO, USA)
30 and Gly-Pro-7-amino-4-trifluoromethylcoumarin (Calbiochem, San Diego, CA, USA) were tested. Other substrates tested were H-Ala-Pro-pNA/HCl, H-Arg-Pro-pNA acetate salt, H-Lys-Ala-pNA.2HCl, H-Asp-Pro-pNA, H-Ala-Ala-pNA/HCl, H-Ala-Ala-Pro-pNA/HCl, H-Ala-Ala-Phe-pNA, succinyl-Ala-Pro-pNA, H-
35 Ala-Phe-Pro-pNA and Z-Ala-Pro-p-NA from Bachem

- 23 -

(Switzerland). H-Ala-Pro-4-methoxy β NA/HCl, Z-Lys-Pro-4-methoxy β NAformate salt, H-Lys-Pro-4-methoxy β NA/HCl, Z-Ala-Pro-4-methoxy β NA, H-Gly-Pro- β NA and H-His-Ser-4-methoxy β NAacetate salt (Bachem) were tested for their
5 ability to stain unfixed transfected cells.
All inhibitors were (see Table 2) incubated with each purified enzyme in phosphate buffer, pH 7.4, for 15 min before the addition of substrate. After the addition of 1mM H-Ala-Pro-pNA substrate for purified DPP8 and 1 mM H-
10 Gly-Pro-pNA substrate for purified DPPIV, samples were incubated for 60 min at 37°C. All enzyme assays were performed in triplicate.

Chromosomal localization of DPP8 by Fluorescence in situ
15 Hybridization (FISH) analysis
DPP8 was localized using two different probes, the DPP8 EST and the T8 clone. The probes were nick-translated with biotin-C¹⁴-dATP and hybridized in situ at a final concentration of 10ng/ul to metaphases from two normal
20 males. The FISH method was modified from that previously described [37] in that chromosomes were stained before analysis with both propidium iodide (as counterstain) and DAPI (for chromosomal identification). Images of metaphase preparations were captured by a cooled CCD camera using
25 the Cyto Vision Ultra image collection and enhancement system (Applied Imaging International Ltd). FISH signals and the DAPI banding pattern were merged for figure preparation.

30 Expression of DPP8 in human lymphocytes and cell lines
RNA (1 μ g) was reverse-transcribed using the Superscript II enzyme kit (Gibco-BRL) as described previously [42]. PCR using DPP8-pr18 (CTGTGACGCCACTAATTATCTATG) as the forward primer and DPP8-pr26R (CCTAGAGAGGCTAGGGTATTCAAG) as the

- 24 -

reverse primer was used to detect full-length DPP8 mRNA. The glyceraldehyde-3-phosphate dehydrogenase (G3PDH) control primer set was G3PDH for (ACCACAGTCCATGCCATCAC) and G3PDHrev (TCCACCACCCTGTTGCTGTA) to give a 470-bp product.

cDNA (diluted 1 : 4; 1µg) was amplified in a 25-µL PCR mixture which contained: 0.2 mM dNTPs, 0.125 unit Amplitaq Gold enzyme (Perkin-Elmer), 1 x buffer II (Perkin-Elmer), 1.5 mM MgCl₂ and 100ng mL⁻¹ each primer. The 35-cycle PCR was performed as follows: denaturation at 94°C for 1 min, primer annealing at 55°C for 30 s, and an extension step at 72°C for 1 min. The amplified products were analyzed by electrophoresis of 15µL PCR mixture on a 3 : 1 Nusieve gel (FMC Bioproducts, Rockville, MD, USA) plus 0.5 µg mL⁻¹ ethidium bromide in Tris/acetate/EDTA buffer (0.04 M Tris/acetate, 0.001 M EDTA, pH 8.0).

Anti-peptide antibody

Methods followed are described in Current Protocols in Immunology [35]. Two peptides were chosen using the software MacVector to predict antigenicity. The two peptides were custom synthesized (Auspep, Melbourne) and conjugated to diphtheria toxin (Auspep, Melbourne). Rabbits were immunized with both peptides and serum collected at time zero and after each injection (IMVS, Adelaide).

The two peptides used were:

PEPTIDE Name: TEDDA-N
SEQUENCE: CTGYTERYMGHPDQNEQG-NH₂

This is amino acids 773 to 789, plus a Cys at the N-

- 25 -

terminus.

PEPTIDE Name: TEDDR-C

SEQUENCE: GKPYDLQIYPQERHSC-NH₂

5

This is amino acids 836 to 850, plus a Cys at the C-terminus.

These sequences were taken from the C-terminal portion of
10 DPP8.

Monoclonal antibody to DPP8

Standard methods were used for antibody production [35].
Mice were immunized with 2×10^7 live COS-7 (African Green
15 Monkey Kidney) cells that had been transiently transfected
with the DPP8 cDNA in the pcDNA3 vector. The final
immunisation was with CHO (Chinese Hamster Ovary) cells
stably transfected with DPP8 cDNA in the pEE14 vector.
Spleen cells were fused with a standard fusion partner,
20 X63Ag8 myeloma cells. Hybridoma culture supernatants were
tested by immunoperoxidase histochemistry on monolayers of
the DPP8-transfected CHO cell line, using untransfected
CHO cells as the negative control. Hybridomas that
produced antibody activity were cloned.

25

RESULTS

Molecular cloning and sequence analysis of DPP8

The insert in ATCC EST AA417787 was 795 bp in length,
containing 527 bp of coding sequence, a TAA stop codon and
30 258 bp of 3' noncoding sequence (Figure 1).

The hybridization of the Master RNA blot revealed that the
gene comprising ESTAA417787 has ubiquitous tissue
expression, with high levels of expression in testis and
35 placenta. Based on this expression pattern, a placental

- 26 -

cDNA library was screened with a 484 bp PCR product produced by the forward and reverse DPP8 primers. Sequence homology analysis revealed that only 2 of 23 clones contained 5' sequence additional to the sequence of ESTAA417787. These cDNA clones were designated T8 and T21, and were 1669 bp and 1197 bp respectively (Figure 1). In addition, comparison of these sequences to ESTAA417787 revealed that T8 cDNA lacked a 153 bp (51aa) region that was present in T21 cDNA and ESTAA417787. This deletion would result in the loss of the catalytic serine (GWSYGG) in T8 cDNA. Many of the other clones characterized appeared to contain unrelated sequence which are probably intronic sequences as a result of incomplete splicing.

The 5' RACE technique was utilized on both ATC RNA and placental RNA to obtain the 5' end of the DPP8 gene. The RACE product obtained from activated T cell RNA was 0.2 kb larger than that from placental RNA but otherwise identical (Figure 1). The first methionine within a Kozak sequence was found 214 bp from the 5' end of the activated T cell RACE product. This 5' 211bp region was 70.5 % GC rich and contained a number of potential promoter and enhancer elements (Sp1, Ap1 and ETF sites) and so was deduced to be the 5' flanking region of the DPP8 gene. In order to confirm the identity of the 5' RACE product as the 5' end of DPP8, RT-PCR was carried out to span across the junction between the RACE product and T8 cDNA library clone. The RT-PCR on ATC RNA produced two clones ATCd3-2-1 and ATC3-3-10 (Figure 1). Compared to T8 and T21, both clones had an additional insert region of 144bp (48 aa) immediately adjacent to the splice site of T8. Sequence homology analysis of this additional insert region found a homologous region in both the *C. elegans* homologue and DPP4. This clearly showed that T8 and T21 library clones represented splice variants of DPP8. The smaller clone

- 27 -

ATCd3-3-10 was also found to represent another splice variant of DPP8 as it contained a 516 bp deletion at the 5' end which would result in a deletion of 175 aa.

5 A full-length DPP8 clone was created using the larger RACE product, ATC3-2-1 and the T21 library clone. This generated a putative DPP8 cDNA of 3.1 kb (including 5' and 3' untranslated regions) with an open reading frame of 882 aa for further sequence analysis and examining DPP8
10 function. This 882 putative DPP8 protein contained no N-linked glycosylation sites and Kyte-Doolittle hydrophobicity analyses revealed it lacked a transmembrane domain, unlike DPP4, FAP and DPP6. Thus it is likely that DPP8 is a cytoplasmic protein (Figure 2). The predicted
15 DPP8 protein shared 51 % amino acid similarity and 27 % amino acid identity with human DPP4; the C termini of these proteins exhibited the most homology (Figure 3).

Tissue distribution of DPP8 as determined by Master RNA and Northern Blot

20 A master RNA blot was probed with a 484 nt PCR product produced by the forward and reverse DPP8 primers as mentioned previously. The mRNA tissue expression of DPP8 was ubiquitous in all human adult and fetal tissues. A
25 similar ubiquitous expression pattern was observed using DPP4 cDNA as a probe (data not shown). However, by visual assessment the greatest levels of expression using each gene specific probe were in different tissues. The most intense signals using the DPP8 probe were in testis
30 followed by placenta whereas the most intense signals using the DPP4 probe were in salivary gland and prostate gland followed by placenta (data not shown). The probes did not bind any of the negative controls on the blot.

35 Northern blot analysis was performed on mRNA derived from

- 28 -

different human tissues (Figure 4). Two DPP8 specific probes indicated the presence of transcripts in all tissues examined. A transcript approximately 3.0 kb in size consistent with the approximate expected size of DPP8 message was detected only in the testis. However, two transcripts of 8.0 and 5.0 kb respectively were present in testis, spleen, peripheral blood leukocytes and ovary at high levels; in prostrate, small intestine, and colonic mucosa at moderate levels; and in the thymus at lower levels. The Multiple tissue Northern blot was also probed with radiolabeled human β -actin probe and a common 2.0 kb transcript was seen in all tissues (Figure 4).

Expression of DPP8 in mice determined by Northern Blot and rtPCR.

The human DPP8 cDNA sequence cross-hybridized with murine derived liver RNA. The Northern blot containing total RNA from mouse liver hybridized to a human DPP8 probe, showing that DPP8 mRNA is expressed in mouse liver (Figure 9A). Two mRNA transcripts of murine DPP8 were present. This is a similar pattern to that observed for human DPP8. These transcripts probably represent different length 5' and 3' untranslated regions of the murine DPP8 gene. The presence of DPP8 mRNA in the mouse liver was also demonstrated using rt-PCR. The primers tested generated a 537bp PCR product. A Southern blot of this product confirmed that the murine DPP8 cross-hybridizes with human DPP8 (Figure 9B).

Expression and functional activity of DPP8

To assess the function of DPP8 protein, the full length DPP8 cDNA of 3.1 kb was cloned into the Xba I site of pcDNA3.1A/V5/His expression vector to produce two constructs. The first construct, pcDNA3.1-DPP8, expressed DPP8 protein on its own whilst the second construct,

- 29 -

pcDNA3.1-DPP8/V5/His expressed a protein with the V5 epitope and His tag fused to the C-terminus of DPP8 to facilitate analysis of protein expression. Mammalian expression constructs were stably transfected into COS-7
5 cells and cellular sonicates prepared. Consistent with the molecular weight predicted from the amino acid sequence a 100 kDa monomer was detected by Western blotting of stable DPP8/V5/His expressing cells (Figure 6). DPP8/V5/His protein was detected in the cytoplasmic compartment but
10 not on the surface of ethanol fixed stable DPP8/V5/His expressing COS cells, using the anti-V5 mAb.

DPP8 is a dipeptidyl peptidase

Sequence homology between DPPIV and DPP8 suggested
15 functional similarities, so cell lysates of DPP8-transfected cells were examined for proline-specific peptidase activity. DPPIV expressed in COS-7 cells with or without the V5/His tag were positive controls, and negative controls included vector-only transfected COS-7
20 cells. Extracts of DPP8-transfected COS-7 cells hydrolyzed H-Ala-Pro-pNA and H-Arg-Pro-pNA but not H-Gly-Pro-pNA, H-Gly-Arg-pNA, H-Gly-Pro-toluenesulfonate or H-Gly-Pro-7-amino-4-trifluoromethylcoumarin above the levels exhibited by untransfected COS-7 cells (data not shown).
25 The pH optimum of DPP8 enzyme activity was 7.4 (Fig. 5A), similar to the pH 7.8 optimum DPPIV enzyme activity [43,44]. DPP8 exhibited little activity below pH 6.3, suggesting that it is not an enzyme of the lysosome/endosome compartment. Of all the substrates
30 tested on cell monolayers, only Ala-Pro-4MβNA/HCl stained DPP8-transfected COS cells and CHO cells (data not shown).

Both purified recombinant DPP8/V5/His and purified recombinant DPPIV/V5/His hydrolyzed H-Ala-Pro-pNA, G-Gly-Pro-pNA and H-Arg-Pro-pNA. Transfection with DPP8
35

- 30 -

possibly causes increased dipeptidase, tripeptidase and endopeptidase activities, similar to an effect of DPPIV transfection of melanoma cells [18]. Indeed, our results showed that DPP8 transfected COS-7 cells, but not purified recombinant DPP8, exhibited tripeptidyl peptidase activity using the substrate H-Ala-Ala-Pro-pNA and endopeptidase activity using the substrate Z-Ala-Pro-pNA (data not shown). This was investigated further, and neither of the tripeptidyl peptidase substrates H-Ala-Ala-Phe-pNA or H-Ala-Phe-Pro-pNA [45] nor the prolyl endopeptidase substrates Z-Ala-Pro-pNA or succinyl-Ala-Pro-pNA were cleaved by purified DPP8. Our data clearly demonstrate that DPP8 is a dipeptidyl peptidase and lacks tripeptidyl peptidase or endopeptidase activities.

The nature of the catalytic mechanism of DPP8 was further investigated using various inhibitors. DPP8 enzyme activity was significantly inhibited by serine proteinase inhibitors and was insensitive to inhibitors of metalloproteinases, aspartyl proteinases and cysteine proteinases. DPP8 enzyme activity was significantly inhibited by zinc, which completely inhibits DPPIV enzyme activity [46]. The peptides Ala-Pro-Gly and Lys-Pro mimic DPP8 substrates and probably competitively inhibited DPP8.

Chromosomal localization of DPP8

Two probes were used for FISH analysis, ESTAA417787 and the T8 clone from the placental library. Seventeen metaphases from the first normal male were examined for fluorescent signal. All of these metaphases showed signal on one or both chromatids of 15 at band q22 (Figure 5). There were a total of 2 non-specific background dots observed in these metaphases. A similar result was obtained from the hybridization of the probe to 15 metaphases from the second normal male (data not shown).

- 31 -

Analysis of DPP8 gene expression by RT-PCR

DPPIV is expressed by most lymphocytes and lymphocytic cell lines but upregulated on activated lymphocytes [47, 41, 48, 49]. The various splice variants of DPP8 might not encode functional protein, so the PCR was designed to detect only mRNA that contained full-length sequence (Fig. 1). At 35 cycles, amplification product of the expected size (783 bp) was readily observed in OKT3-stimulated PBMCs (six of six subjects; Fig. 8) but not in unstimulated PBMCs from most subjects (four of five, Fig. 8A), suggesting that more DPP8 mRNA is expressed in activated T cells than in unstimulated PBMCs. Similar RT-PCR data were obtained from PBMCs stimulated with phytohaemagglutinin (data not shown). In addition, DPP8 mRNA was expressed in all B and T cell lines examined and in both liver and placenta (Fig. 8B).

Anti-peptide antibody

The sera of two rabbits were tested by ELISA in peptide-coated wells. Both sera bound both peptides whereas the pre-immunisation serum samples did not exhibit specific binding. Western blots on extracts of cell lines, cell lines transfected with DPP8 cDNA and activated human lymphocytes showed that a rabbit antiserum to the two DPP8 peptides binds a 100kDa band, which is the size of DPP8. (Data not shown).

Table 1. K_m and V_{max} values for DPP8 and DPPIV

	K_m (mM)		V_{max} ($\Delta A \text{ min}^{-1} \times 1000$)	
	DPPIV	DPP8	DPPIV	DPP8
H-Ala-Pro-pNA	0.374 ± 0.134	0.991 ± 0.171	9.6 ± 1.0	12.4 ± 0.9
H-Gly-Pro-pNA	0.347 ± 0.088	0.467 ± 0.064	7.2 ± 0.49	3.5 ± 0.14

5

Table 2. Inhibition of the peptidase activity of DPP8 in
 10 comparison with DPPIV. Common proteinase inhibitors of
 various enzyme types were incubated with the purified
 peptidases before assay with the substrates H-Ala-Pro-pNA
 on DPP8 or H-Gly-Pro-pNA on DPPIV. AEBSF, 4-(2-
 aminoethyl)benzenesulfonylfluoride.

15

Type of inhibitor	Concentration	Residual activity (% of control)	
		DPP8	DPPIV
None		100	100
Serine proteinase			
AEBSF	4 mM	40	52
Aprotinin	4 $\mu\text{g mL}^{-1}$	47	81
Benzamidine/HCl	10 mM	82	89
Peptides			
Gly-Gly-Gly	10 mM	99	106
Ala-Pro-Gly	10 mM	51	67
H-Lys-Pro-OH HCl salt	4 mM	63	45
Zinc sulphate	2 mM	25	0
Metalloproteinase			
EDTA	2 mM	115	99
Aspartate (acidic) proteinase			
Pepstatin	2 $\mu\text{g mL}^{-1}$	107	110
Leupeptin	0.1 mM	93	104
Cysteine (thiol) proteinase			
Iodoacetamide	2 mM	100	115
Dithiothreitol	2 mM	108	109

- 33 -

Discussion

We describe the cloning, recombinant expression, biochemistry and tissue expression of a novel human DPPIV-related postproline peptidase that we have named DPP8. DPP8 exhibited dipeptidyl aminopeptidase but not tripeptidyl peptidase or endopeptidase activity. Like DPPIV, DPP8 was found to exhibit significant mRNA expression in activated T cells. Clear indications that DPP8 is a monomeric, nonglycosylated, soluble, cytoplasmic protein, which are characteristics of PEP but not of DPPIV, FAP or DPP6, were provided by our sequence and localisation data. DPP8 enzyme activity had a neutral pH optimum, suggesting that it is not active in the acidic lysosome/endosome compartment.

By homology with DPPIV, DPP8 is a member of the DPPIV-like gene family, a member of the prolyl oligopeptidase family S9b, and a member of the enzyme clan SC. The residues in DPP8 that potentially form the charge-relay system are Ser739, Asp817 and His849 (Fig.2). The dipeptidyl peptidase activity of DPP8 and the absence of detectable tripeptidyl peptidase or endopeptidase activities by purified DPP8 further support its placement in the S9b family. Furthermore, the DPP8 substrate specificity was indistinguishable from that of the structurally related peptidases DPPIV and FAP.

The role of DPPIV in human lymphocytes has been studied in detail using enzyme inhibitors [49, 50-54]. DPPIV-specific inhibitors suppress both DNA synthesis and cytokine production in vitro [48, 49, 52]. In addition, DPPIV-specific inhibitors decrease phorbol myristate acetate-induced tyrosine phosphorylation in human lymphocytes, further suggesting a role for DPPIV enzyme

- 34 -

activity in lymphocyte activation [54]. *In vivo*, inhibitors of DPPIV suppress arthritis [20] and prolong cardiac allograft survival in animal models [55]. The ability of DPP8 to cleave DPPIV substrates indicates that DPPIV inhibitors may also inhibit DPP8 and that inhibitor studies may require further interpretation. Indeed, DPP8 may be responsible for some of the physiological functions that have been assigned to DPPIV.

FAP and DPPIV are integral membrane glycoproteins and require dimerization for catalytic activity [9, 56, 57]. In contrast, DPP8 and PEP are non-glycosylated cytosolic proteins that are catalytically active as monomers [58] and cleave Pro-Xaa bonds [43, 59]. However, the substrate specificity of PEP is distinct from DPP8. PEP is an endopeptidase that does not cleave if a free α -amine lies N-terminal to the proline (e.g. it does not cleave H-Ala-Pro). Recently we have proposed that the tertiary structure of DPPIV is similar to that of PEP in having a seven-blade β -propeller domain and an α/β -hydrolase domain [3, 39, 1]. The significant sequence identity between DPP8 and DPPIV indicates that the tertiary structures of DPP8 and DPPIV are similar. However, DPP8 contains 110 amino acids more than DPPIV, so it could have an additional element of tertiary structure such as an eighth propeller blade.

The ancestral relationships between DPP8, DPPIV and FAP are reflected in their chromosomal localization. While DPPIV and FAP have both been localized to the long arm of chromosome 2, 2q24.3 [60] and 2q23 [61] respectively, DPP8 was localized to 15q22. The related genes DPP6 and PEP have been localized to chromosome 7 [62] and 6q22 respectively [63].

35

- 35 -

Two human disease loci have been mapped to 15q22. These loci are an autosomal recessive deafness locus [64] and a form of Bardet-Biedl syndrome, type 4 [65]. Two of the clinical manifestations of Bardet-Biedl syndrome are obesity and diabetes. Attractin [66] and DPPIV have roles in obesity [67] and diabetes [22, 68, 15] respectively and as their substrate specificities overlap with that of DPP8, it is possible that DPP8 may be involved in Bardet-Biedl syndrome.

DPPIV is expressed on the surface of T cells and is a costimulatory molecule called CD26 [3]. CD26-negative cell lines have residual DPPIV enzyme activity and PBMC have non-DPPIV derived activity against Ala-Pro substrates [69], indicating the existence of other peptidase(s) with DPPIV-like activity. DPPIV- β exhibits a peptidase activity similar to DPPIV but is a 70-80 kDa cell surface glycoprotein [70] and is therefore distinct from DPP8.

The biological significance of the three splice variants of DPP8 that we discovered is unknown. None of these splice variants result in a frame shift or premature protein termination (Fig. 1). Two of the splice variants contain all the predicted catalytic triad residues and thus potentially produce proteins with peptidase activity. Alternate splice forms of FAP mRNA have also been observed [71, 72]. It is possible that expression of splice variants may be used to regulate the levels of active protein. DPP8 Northern blots revealed a number of differently sized transcripts. The predicted sizes of splice variants of DPP8 ranged from 2.6 to 3.1 kb whereas the large transcripts seen in most tissues examined in the Northern blots were 8.5 kb and 5.0 kb respectively. Similarly, two other members of the DPPIV-like gene family, DPPIV and DPP6, exhibit mRNA transcripts in

- 36 -

Northern blots that are much larger than the cDNA size [60, 61]. We propose that the major transcripts for DPP8 mRNA and its splice variants lie within the 5 kb band while the 8.5 kb transcript(s) may contain additional 5' and 3' untranslated sequences. DPP8 appears to be like DPPIV in having a ubiquitous mRNA expression pattern by Northern analysis while being upregulated in activated T cells. The similarities between DPP8 and DPPIV suggest that DPP8 may, like DPPIV, play a role in T cell costimulation and proliferation. The development of DPP8 specific antibodies or inhibitors will facilitate work in this area.

In summary, we have identified and characterized a novel human dipeptidyl aminopeptidase DPP8 with structural and functional similarities to DPPIV and FAP. With many diverse biological roles suggested for DPPIV, particularly in the immune system, and the roles of FAP in tumor growth and liver disease, it will be interesting to investigate the roles of this new member of the DPPIV-like gene family in these systems. Further work in understanding this novel protein and the elucidation of inhibitors and physiological substrates will help identify the specific functions of individual members of this gene family.

- 37 -

REFERENCES

1. Abbott CA, GW McCaughan & MD Gorrell 1999 Two highly conserved glutamic acid residues in the predicted beta propeller domain of dipeptidyl peptidase IV are required for its enzyme activity *FEBS Letters* 458: 278-84.
2. Abbott CA, DMT Yu, GW McCaughan & MD Gorrell 2000 Post proline peptidases having DP IV like enzyme activity *Advances in Experimental Medicine and Biology* 477: 103-9.
3. McCaughan GW, MD Gorrell, GA Bishop, CA Abbott, NA Shackel, PH McGuinness, MT Levy, AF Sharland, DG Bowen, D Yu, L Slaitini, WB Church & J Napoli 2000 Molecular pathogenesis of liver disease: an approach to hepatic inflammation, cirrhosis and liver transplant tolerance *Immunological Reviews* 174: 172-91.
4. Scanlan MJ, BK Raj, B Calvo, P Garin-Chesa, MP Sanz-Moncasi, JH Healey, LJ Old & WJ Rettig 1994 Molecular cloning of fibroblast activation protein alpha, a member of the serine protease family selectively expressed in stromal fibroblasts of epithelial cancers *Proceedings of the National Academy of Sciences United States of America* 91: 5657-61.
5. Handbook of Proteolytic Enzymes.. Barrett AJ, ND Rawlings & JF Woess. 1998., London: Academic Press. 1666.
6. Jacotot E, C Callebaut, J Blanco, B Krust, K Neubert, A Barth & AG Hovanessian 1996 Dipeptidyl-peptidase IV-beta, a novel form of cell-surface-expressed protein with dipeptidyl-peptidase IV activity *European Journal of Biochemistry* 239: 248-58.
7. Rawlings ND & AJ Barrett 1999 MEROPS: the peptidase database *Nucleic Acids Research* 27: 325-31.
8. Park JE, MC Lenter, RN Zimmermann, P Garin-Chesa, LJ Old & WJ Rettig 1999 Fibroblast activation protein: A dual-specificity serine protease expressed in reactive human tumor stromal fibroblasts *Journal of*

- 38 -

Biological Chemistry 274: 36505-12.

9. Levy MT, GW McCaughan, CA Abbott, JE Park, AM Cunningham, E Muller, WJ Rettig & MD Gorrell 1999 Fibroblast activation protein: A cell surface dipeptidyl
5 peptidase and gelatinase expressed by stellate cells at the tissue remodelling interface in human cirrhosis *Hepatology* 29: 1768-78.
10. De Meester I, S Korom, J Van Damme & S Scharpé 1999 CD26, let it cut or cut it down *Immunology Today* 20:
10 367-75.
11. Natural substrates of dipeptidyl peptidase IV. De Meester I, C Durinx, G Bal, P Proost, S Struyf, F Goossens, K Augustyns & S Scharpé. 2000, in *Cellular Peptidases in Immune Functions and Diseases II*, J Langner
15 & S Ansorge, Editor. Kluwer, New York. p. 67-88.
12. Mentlein R 1999 Dipeptidyl-peptidase IV (CD26): role in the inactivation of regulatory peptides *Regulatory Peptides* 85: 9-24.
13. Morrison ME, S Vijayasaradhi, D Engelstein, AP
20 Albino & AN Houghton 1993 A marker for neoplastic progression of human melanocytes is a cell surface ectopeptidase *Journal of Experimental Medicine* 177: 1135-43.
14. Mueller SC, G Gherzi, SK Akiyama, QXA Sang, L
25 Howard, M Pineiro-Sanchez, H Nakahara, Y Yeh & WT Chen 1999 A novel protease-docking function of integrin at invadopodia *Journal of Biological Chemistry* 274: 24947-52.
15. Holst JJ & CF Deacon 1998 Inhibition of the activity of dipeptidyl-peptidase IV as a treatment for
30 type 2 diabetes *Diabetes* 47: 1663-70.
16. Marguet D, L Baggio, T Kobayashi, AM Bernard, M Pierres, PF Nielsen, U Ribel, T Watanabe, DJ Drucker & N Wagtman 2000 Enhanced insulin secretion and improved glucose tolerance in mice lacking CD26 *Proceedings of the*

- 39 -

National Academy of Sciences of the United States of America 97: 6874-9.

17. Ohtsuki T, H Tsuda & C Morimoto 2000 Good or evil: CD26 and HIV infection *Journal of Dermatological Science* 22: 152-60.
18. Wesley UV, AP Albino, S Tiwari & AN Houghton 1999 A role for dipeptidyl peptidase IV in suppressing the malignant phenotype of melanocytic cells *Journal of Experimental Medicine* 190: 311-22.
19. Korom S, I De Meester, THW Stadlbauer, A Chandraker, M Schaub, MH Sayegh, A Belyaev, A Haemers, S Scharpé & JW Kupiecweglinski 1997 Inhibition of CD26/dipeptidyl peptidase IV activity in vivo prolongs cardiac allograft survival in rat recipients *Transplantation* 63: 1495-500.
20. Tanaka S, T Murakami, H Horikawa, M Sugiura, K Kawashima & T Sugita 1997 Suppression of arthritis by the inhibitors of dipeptidyl peptidase IV *International Journal of Immunopharmacology* 19: 15-24.
21. Augustyns K, G Bal, G Thonus, A Belyaev, XM Zhang, W Bollaert, AM Lambeir, G Durinx, F Goossens & A Haemers 1999 The unique properties of dipeptidyl-peptidase IV (DPP IV / CD26) and the therapeutic potential of DPP IV inhibitors *Current Medicinal Chemistry* 6: 311-27.
22. Hinke SA, JA Pospisilik, HU Demuth, S Mannhart, K Kuhn-Wache, T Hoffmann, E Nishimura, RA Pederson & CHS McIntosh 2000 Dipeptidyl peptidase IV (DPIV/CD26) degradation of glucagon - Characterization of glucagon degradation products and DPIV-resistant analogs *Journal of Biological Chemistry* 275: 3827-34.
23. Korom S, I De Meester, A Coito, E Graser, HD Volk, K Schwemmler, S Scharpe & JW Kupiec-Weglinski 1999 Immunomodulatory influence of CD26 dipeptidylpeptidase IV during acute and accelerated rejection *Langenbecks Archives of Surgery* 1: 241-5.

- 40 -

24. Tavares W, DJ Drucker & PL Brubaker 2000
Enzymatic- and renal-dependent catabolism of the
intestinotropic hormone glucagon-like peptide-2 in rats
American Journal of Physiology Endocrinology and
5 *Metabolism* 278: E134-E9.
25. David F, AM Bernard, M Pierres & D Marguet 1993
Identification of serine 624, aspartic acid 702, and
histidine 734 as the catalytic triad residues of mouse
dipeptidyl-peptidase IV (CD26). A member of a novel family
10 of nonclassical serine hydrolases *J Biol Chem* 268: 17247-
52.
26. Ogata S, Y Misumi, E Tsuji, N Takami, K Oda & Y
Ikehara 1992 Identification of the active site residues in
dipeptidyl peptidase IV by affinity labeling and site-
15 directed mutagenesis *Biochemistry* 31: 2582-7.
27. Dipeptidyl peptidase IV (DPPIV/CD26):
biochemistry and control of cell-surface expression.
Trugnan G, T Ait-Slimane, F David, L Baricault, T Berbar,
C Lenoir & C Sapin. 1997, in *Cell-Surface Peptidases in*
20 *Health and Disease*, AJ Kenny & CM Boustead, Editor. BIOS
Scientific Publishers: Oxford. pp 203-17.
28. Steeg CS, U Hartwig & B Fleischer 1995 Unchanged
signaling capacity of mutant CD26/dipeptidylpeptidase IV
molecules devoid of enzymatic activity *Cell Immunol* 164:
25 311-5.
29. Fülöp V, Z Böcskei & L Polgar 1998 Prolyl
oligopeptidase - an unusual beta-propeller domain
regulates proteolysis *Cell* 94: 161-70.
30. Ausubel FM, R Brent, RE Kingston, DD Moore, JG
Seidman, JA Smith & K Struhl, ed. *Current Protocols in*
30 *Molecular Biology*. 1998, John Wiley & Sons: USA.
31. Molecular cloning: a laboratory manual.
Sambrook J, EF Fritsch & T Maniatis. 1989. 2nd ed., Cold
Spring Harbor: Cold Spring Harbor Laboratory Press.

- 41 -

32. Augustyns KJL, AM Lambeir, M Borloo, I Demeester, I Vedernikova, G Vanhoof, D Hendriks, S Scharpe & A Haemers 1997 Pyrrolidides - synthesis and structure-activity relationship as inhibitors of dipeptidyl
5 peptidase IV *European Journal of Medicinal Chemistry* 32: 301-9.
33. Stockel-Maschek A, C Mrestani-Klaus, B Stiebitz, HU Demuth & K Neubert 2000 Thioxo amino acid pyrrolidides and thiazolidides: new inhibitors of proline
10 specific peptidases *Biochimica et Biophysica Acta - Protein Structure & Molecular Enzymology* 1479: 15-31.
34. Schön E, I Born, HU Demuth, J Faust, K Neubert, T Steinmetzer, A Barth & S Ansorge 1991 Dipeptidyl
15 peptidase IV in the immune system. Effects of specific enzyme inhibitors on activity of dipeptidyl peptidase IV and proliferation of human lymphocytes *Biological Chemistry Hoppe Seyler* 372: 305-11.
35. Coligan JE, AM Kruisbeek, DH Margulies, EM Shevach & W Strober, eds. Current Protocols in Immunology.
20 1998, John Wiley & Sons: USA.
36. Fibroblast activation protein. Rettig WJ. 1998, in *Handbook of Proteolytic Enzymes*, AJ Barrett, ND Rawlings & JF Woessner, Editor. Academic Press: San Diego. p. 387-9.
- 25 37. McCaughan, G. W., Siah, C. L., Abbott, C., Wickson, J., Ballesteros, M. & Bishop, G. A. (1993) Dipeptidyl peptidase IV is down-regulated in rat hepatoma cells at the mRNA level, *J. Gastroenterol. Hepatol.* 8, 142-5.
- 30 38. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. (1990) Basic local alignment search tool, *J. Mol. Biol.* 215, 403-410.
33. Callen, D. F., Baker, E., Eyre, H. J., Chernos, J. E., Bell, J. A. & Sutherland, G. R. (1990) Reassessment of
35 two apparent deletions of chromosome 16p to an ins(11;16)

- 42 -

and a t(1;16) by chromosome painting, *Annales De Genetique*. 33, 219-21.

39. Abbott, C. A., McCaughan, G. W., Levy, M. T., Church, W. B. & Gorrell, M. D. (1999) Binding to human
5 dipeptidyl peptidase IV by adenosine deaminase and antibodies that inhibit ligand binding involves overlapping, discontinuous sites on a predicted beta propeller domain, *Eur. J. Biochem*. 266, 798-810.

40. McCaughan, G. W., Wickson, J. E., Creswick, P.
10 F. & Gorrell, M. D. (1990) Identification of the bile canalicular cell surface molecule GP110 as the ectopeptidase dipeptidyl peptidase IV: an analysis by tissue distribution, purification and N-terminal amino acid sequence, *Hepatology*. 11, 534-44.

15 41. Gorrell, M. D., Wickson, J. & McCaughan, G. W. (1991) Expression of the rat CD26 antigen (dipeptidyl peptidase IV) on subpopulations of rat lymphocytes, *Cell. Immunol*. 134, 205-215.

42. Bishop, G. A., Rokahr, K. L., Napoli, J. &
20 McCaughan, G. W. (1993) Intragraft cytokine mRNA levels in human liver allograft rejection analysed by reverse transcription and semiquantitative polymerase chain reaction amplification, *Transplant Immunol*. 1, 253-61.

43. Yoshimoto, T., Fischl, M., Orlowski, R. C. &
25 Walter, R. (1978) Post-proline cleaving enzyme and post-proline dipeptidylpeptidase, *J. Biol. Chem*. 253, 3708-3716.

44. Fukasawa, K. M., Fukasawa, K., Hiraoka, B. Y. & Harada, M. (1981) Comparison of dipeptidyl peptidase IV
30 prepared from pig liver and kidney, *Biochim. Biophys. Acta*. 657, 179-89.

45. Banbula, A., Mak, P., Bugno, M., Silberring, J., Dubin, A., Nelson, D., Travis, J. & Potempa, J. (1999)
Prolyl tripeptidyl peptidase from *Porphyromonas*
35 *gingivalis*. A novel enzyme with possible pathological

- 43 -

implications for the development of periodontitis, *J. Biol. Chem.* 274, 9246-9252.

46. Wolf, G. B., Scherberich, J. E., Fischer, P. & Schoeppe, W. (1989) Isolation and characterization of
5 dipeptidyl aminopeptidase IV from human kidney cortex, *Clin. Chim. Acta.* 179, 61-71.

47. Fox, D. A., Hussey, R. E., Fitzgerald, K. A., Acuto, O., Poole, C., Palley, L., Daley, J. F., Schlossman, S. F. & Reinherz, E. L. (1984) Tal, a novel
10 105 KD human T cell activation antigen defined by a monoclonal antibody, *J. Immunol.* 133, 1250-6.

48. Bühling, F., Kunz, D., Reinhold, D., Ulmer, A. J., Ernst, M., Flad, H. D. & Ansorge, S. (1994) Expression and functional role of dipeptidyl peptidase IV (CD26) on
15 human natural killer cells, *Natural Immunity.* 13, 270-9.

49. Bühling, F., Junker, U., Reinhold, D., Neubert, K., Jager, L. & Ansorge, S. (1995) Functional role of CD26 on human B lymphocytes, *Immunol Lett.* 45, 47-51.

50. Flentke, G. R., Munoz, E., Huber, B. T., Plaut, A. G., Kettner, C. A. & Bachovchin, W. W. (1991) Inhibition of dipeptidyl aminopeptidase IV (DP-IV) by Xaa-boroPro dipeptides and use of these inhibitors to examine
20 the role of DP-IV in T-cell function, *Proc. Natl. Acad. Sci. USA.* 88, 1556-9.

51. Schön, E., Jahn, S., Kiessig, S. T., Demuth, H. U., Neubert, K., Barth, A., Von-Bähr, R. & Ansorge, S. (1987) The role of dipeptidyl peptidase IV in human T lymphocyte activation. Inhibitors and antibodies against dipeptidyl peptidase IV suppress lymphocyte proliferation
25 and immunoglobulin synthesis in vitro, *Eur. J. Immunol.* 17, 1821-6.

52. Schön, E., Born, I., Demuth, H. U., Faust, J., Neubert, K., Steinmetzer, T., Barth, A. & Ansorge, S. (1991) Dipeptidyl peptidase IV in the immune system.
35 Effects of specific enzyme inhibitors on activity of

- 44 -

dipeptidyl peptidase IV and proliferation of human lymphocytes, *Biol. Chem. Hoppe Seyler.* 372, 305-11.

53. Reinhold, D., Hemmer, B., Gran, B., Born, I., Faust, J., Neubert, K., McFarland, H. F., Martin, R. & Ansorge, S. (1998) Inhibitors of dipeptidyl peptidase IV/CD26 suppress activation of human MBP-specific CD4+ T cell clones, *J. Neuroimmunol.* 87, 203-9.

54. Kähne, T., Neubert, K., Faust, J. & Ansorge, S. (1998) Early phosphorylation events induced by DPIV/CD26-specific inhibitors, *Cell. Immunol.* 189, 60-66.

55. Korom, S., De Meester, I., Stadlbauer, T. H. W., Chandraker, A., Schaub, M., Sayegh, M. H., Belyaev, A., Haemers, A., Scharpé, S. & Kupiecweglinski, J. W. (1997) Inhibition of CD26/dipeptidyl peptidase IV activity in vivo prolongs cardiac allograft survival in rat recipients, *Transplantation.* 63, 1495-1500.

56. Bednarczyk, J. L., Carroll, S. M., Marin, C. & McIntyre, B. W. (1991) Triggering of the proteinase dipeptidyl peptidase IV (CD26) amplifies human T lymphocyte proliferation, *J. Cell. Biochem.* 46, 206-18.

57. De Meester, I., Vanhoof, G., Hendriks, D., Demuth, H. U. & Yaron, A. & Scharpé, S. (1992) Characterization of dipeptidyl peptidase IV (CD26) from human lymphocytes, *Clin. Chim. Acta.* 210, 23-34.

58. Goossens, F., De Meester, I., Vanhoof, G., Hendriks, D., Vriend, G. & Scharpé, S. (1995) The purification, characterization and analysis of primary and secondary-structure of prolyl oligopeptidase from human lymphocytes. Evidence that the enzyme belongs to the alpha/beta hydrolase fold family, *Eur. J. Biochem.* 233, 432-441.

59. Walter, R., Simmons, W. H. & Yoshimoto, T. (1980) Proline specific endo- and exopeptidases, *Mol. Cell. Bioc.* 30, 111-27.

- 45 -

60. Abbott, C. A., Baker, E., Sutherland, G. R. & McCaughan, G. W. (1994) Genomic organisation, exact localization, and tissue expression of the human CD26 (dipeptidyl peptidase IV) gene, *Immunogenetics*. 40, 331-338.
61. Mathew, S., Scanlan, M. J., Mohan Raj, B. K., Murty, V. V., Garin-Chesa, P., Old, L. J., Rettig, W. J. & Chaganti, R. S. (1995) The gene for fibroblast activation protein alpha (FAP), a putative cell surface-bound serine protease expressed in cancer stroma and wound healing, maps to chromosome band 2q23, *Genomics*. 25, 335-337.
62. Yokotani, N., Doi, K., Wenthold, R. J. & Wada, K. (1993) Non-conservation of a catalytic residue in a dipeptidyl aminopeptidase IV-related protein encoded by a gene on human chromosome 7, *Hum. Mol. Genet.* 2, 1037-9.
63. Goossens, F. J., Wauters, J. G., Vanhoof, G. C., Bossuyt, P. J., Schatteman, K. A., Loens, K. & Scharpé, S. (1996) Subregional mapping of the human lymphocyte prolyl oligopeptidase gene (PREP) to human chromosome 6q22, *Cytogenet Cell Genet.* 74, 99-101.
64. Campbell, D. A., McHale, D. P., Brown, K. A., Moynihan, L. M., Houseman, M., Karbani, G., Parry, G., Janjua, A. H., Newton, V., Al-Gazali, L., Markham, A. F., Lench, N. J. & Mueller, R. F. (1997) A new locus for non-syndromal, autosomal recessive, sensorineural hearing loss (DFNB16) maps to human chromosome 15q21-q22, *J. Med. Genet.* 34, 1015-7.
65. Bruford, E. A., Riise, R., Teague, P. W., Porter, K., Thomson, K. L., Moore, A. T., Jay, M., Warburg, M., Schinzel, A., Tommerup, N., Tornqvist, K., Rosenberg, T., Patton, M., Mansfield, D. C. & Wright, A. F. (1997) Linkage mapping in 29 Bardet-Biedl syndrome families confirms loci in chromosomal regions 11q13, 15q22.3-q23, and 16q21, *Genomics*. 41, 93-9.

- 46 -

66. Duke-Cohan, J. S., Gu, J., McLaughlin, D. F., Xu, Y., Freeman, G. J. & Schlossman, S. F. (1998) Attractin (DPPT-L), a member of the CUB family of cell adhesion and guidance proteins, is secreted by activated
5 human T lymphocytes and modulates immune cell interactions, *Proc. Natl. Acad. Sci. USA* 95, 11336-11341.
67. Gunn, T. M., Miller, K. A., He, L., Hyman, R. V., Davis, R. W., Azarani, A., Schlossman, S. F., Duke-Cohan, J. S. & Barsh, G. S. (1999) The mouse mahogany
10 locus encodes a transmembrane form of human attractin, *Nature* 398, 152-156.
68. Drucker, D. J., Shi, Q., Crivici, A., Summersmith, M., Tavares, W., Hill, M., Deforest, L., Cooper, S. & Brubaker, P. L. (1997) Regulation of the
15 biological activity of glucagon-like peptide 2 in vivo by dipeptidyl peptidase IV, *Nature Biotechnol.* 15, 673-677.
69. Smith, R. E., Reynolds, C. J. & Elder, E. A. (1992) The evolution of proteinase substrates with special reference to dipeptidylpeptidase IV, *Histochem. J.* 24,
20 637-47.
70. Blanco, J., Nguyen, C., Callebaut, C., Jacotot, E., Krust, B., Mazaleyrat, J. P., Wakselman, M. & Hovanessian, A. G. (1998) Dipeptidyl-peptidase IV-beta. Further characterization and comparison to dipeptidyl-
25 peptidase IV activity of CD26, *Eur. J. Biochem.* 256, 369-378.
71. Niedermeyer, J., Scanlan, M. J., Garin-Chesa, P., Daiber, C., Fiebig, H. H., Old, L. J., Rettig, W. J. & Schnapp, A. (1997) Mouse fibroblast activation protein:
30 molecular cloning, alternative splicing and expression in the reactive stroma of epithelial cancers, *Int. J. Cancer* 71, 383-9.
72. Goldstein, L. A. & Chen, W. T. (2000) Identification of an alternatively spliced seprase mRNA

- 47 -

that encodes a novel intracellular isoform, *J. Biol. Chem.* 275, 2554-2559.

73. Hough, R. B., Lengeling, A., Bedian, V., Lo, C. & Bucan, M. (1998) Rump white inversion in the mouse
5 disrupts dipeptidyl aminopeptidase-like protein 6 and causes dysregulation of Kit expression, *Proc. Natl. Acad. Sci. USA.* 95, 13800-13805

- 48 -

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A peptide which comprises:
(a) the sequence shown in SEQ ID NO:1; or
5 (b) the amino acid sequences:
His⁷³⁶GlyTrpSerTyrGlyGlyTyrLeu;
Leu⁸¹⁶AspGluAsnValHisPheAlaHis; Glu⁸⁴⁷ArgHisSerIleArg and
Phe²⁵⁵ValLeuGlnGluGluPhe, and which has the substrate
specificity of the sequence shown in SEQ ID NO:1; or
10 (c) the sequence which has at least 60% identity
with the sequence shown in SEQ ID NO:1, and which has the
substrate specificity of the sequence shown in SEQ ID
NO:1.
- 15 2. A peptide according to claim 1 (c), wherein the
amino acid identity is at least 75%.
3. A peptide according to claim 1 (c) wherein the
amino acid identity is at least 95%.
- 20 4. A fragment of the sequence shown in SEQ ID NO:1
which has the substrate specificity of the sequence shown
in SEQ ID NO:1.
- 25 5. A fragment according to claim 4 which consists
of the sequence shown in SEQ ID NO.s: 3, 5 or 7.
6. A peptide according to claim 1, wherein an
asparagine residue in the peptide is not linked to a
30 carbohydrate molecule.
7. A peptide according to claim 1, wherein the
peptide is not expressed on the cell surface membrane of a
cell.

- 49 -

8. A fusion protein comprising the amino acid sequence shown in SEQ ID NO:1 linked with a further amino acid sequence, the fusion protein having the substrate specificity of the sequence shown in SEQ ID NO:1.

5

9. A fusion protein according to claim 8 wherein the further amino acid sequence is selected from the group consisting of GST, V5 epitope and His tag.

10 10. A method of identifying a molecule capable of inhibiting cleavage of a substrate by DPP8 comprising the following steps:

- (a) contacting DPP8 with the molecule;
- (b) contacting DPP8 of step (a) with a substrate capable of being cleaved by DPP8, in conditions sufficient for cleavage of the substrate by DPP8; and
- (c) detecting substrate not cleaved by DPP8, to identify that the molecule is capable of inhibiting cleavage of the substrate by DPP8.

20

11. A method of identifying a molecule capable of inhibiting specifically, the cleavage of a substrate by DPP8, the method comprising the following steps:

- (a) contacting DPP8 and a further protease with the molecule;
- (b) contacting DPP8 and the further protease of step (a) with a substrate capable of being cleaved by DPP8 and the further protease, in conditions sufficient for cleavage of the substrate by DPP8 and the further protease; and
- (c) detecting substrate not cleaved by DPP8, but cleaved by the further protease, to identify that the molecule is capable of inhibiting specifically, the cleavage of the substrate by DPP8.

35

- 50 -

12. A method of reducing or inhibiting the catalytic activity of DPP8, the method comprising the step of contacting DPP8 with an inhibitor of DPP8 catalytic activity.

5

13. A method of cleaving a substrate comprising the step of contacting the substrate with DPP8 in conditions sufficient for cleavage of the substrate by DPP8.

10 14. A method of detecting an activated T cell, the method comprising the step of measuring the level of DPP8 gene expression in a T cell.

15 15. A method according to claim 14, wherein the level of DPP8 gene expression is detected by detecting the amount of DPP8 RNA in the cell.

16. A nucleic acid molecule which:
(a) encodes the sequence shown in SEQ ID NO:1; or
20 (b) consists of the sequence shown in SEQ ID NO:2; or
(c) is capable of hybridizing to a nucleic acid molecule consisting of the sequence shown in SEQ ID NO:2 in stringent conditions, and which encodes a peptide which has the substrate specificity of the sequence shown in SEQ
25 ID NO:1.

17. A nucleic acid molecule according to claim 16 (c) wherein the molecule is capable of hybridising in high stringent conditions.

30

18. A nucleic acid molecule according to claim 16 which is capable of hybridising to a gene which is located at band q 22 on human chromosome 15.

35 19. A nucleic acid molecule according to claim 16

- 51 -

which does not contain 5' or 3' untranslated regions.

20. A fragment of a nucleic acid molecule
consisting of the sequence shown in SEQ ID NO:2, which
5 encodes a peptide which has the substrate specificity of
the sequence shown in SEQ ID NO:1.

21. A fragment according to claim 20 which consists
of the sequence shown in any one of SEQ ID NO.s: 4, 6 or
10 8.

22. A vector comprising a nucleic acid molecule
according to claim 16.

15 23. A cell comprising a vector according to claim
22.

24. A composition comprising a peptide according to
claim 1.
20

25. An antibody which is capable of binding to a
peptide according to claim 1.

26. An antibody according to claim 25 which is
25 produced by a hybridoma cell.

27. A hybridoma cell capable of making an antibody
according to claim 26.

1/9




FIGURE 1

2/9

1 AAGTCTAAAGCCCTCCGAGGCCAAGGCCCTGCTACTGCGCGCGCTGCTTCTTAGTCCGCGCTTCGCGCGCTGGGTTCACCGCGCGCGCGCGCGAGGA 100
 101 AGCCACTGCAACACGAGGACCGGAGTGGAGGCGCGCGAGCATGAAGCGCGCGCGCGCGCGCTCCATAGCGCACGTCGGGACGCTCCGCGCGCGCGCGCGGGA 200
 201 AGGAAATGCAACATCGGAGGACCGGAGTGGAGGCGCGCGAGCATGAAGCGCGCGCGCGCGCGCTCCATAGCGCACGTCGGGAGGAAATATTGAATCAGGAGTGG 300
 301 CCTAAATGAGCCCTTTTATGTTGAGCGGTATTCCTGGAGTCAGCTTAAAGCTGCTTGGGATACCCAGAAATATCATGGTACATCATGCTTAAGG 400
 30 P K L E P F Y V E R Y S W S Q L K K L L A D T R K Y H G Y H M A K A 63
 401 CACCACATGATTTCATGTTGTAAGAGGAATGATCCAGATCGACCTCATTACAGACAGAACTATTACCTTGGCATGCTGCTGAGAACACAGAAATAC 500
 63 P H D F M F V K R M D P D C P H S D R I Y Y L A H S G E N T 96
 501 ACTGTTTATTTGAAATCCCAAACTATCAATAGAGCAGCAGCTTAAATGCTCTCTTGGAGGCTCTTTTGGATCTTTTACGGCAACACTGCACTAT 600
 96 L F Y S E I P K T I N R A A V L H L S W K P L L D L F Q A T L D Y 129
 601 CGAATGTTATTCGAGAGGAAGAACTATTAGAGAAAGAAACCGCATGGAACAGCTCGGAATGCTTCTTACGATTATCACCAGGAGTGGAAATTTG 700
 130 G H Y S R E E L L R E R K R I G T V G I A S Y D Y H Q G S G T F L 163
 701 TGTTCAGGCGGTAGTGAATTTATCAGTAAAGATGGAGGCGCGAGGATTTACGCAACACCTTTAAGCGCCAACTAGTGGAAATAGTTGCTC 800
 163 F O A G S G I Y H V K D G G P Q G F T Q O P L R P N L V E T S C P 196
 801 CAACATACGGATGGATCCAAATATGCCCCGCTGATCCAGACTGGATGCTTTTATACATACCAAGCATTTGGATATCTAACATCGTAACAGAGAA 900
 196 M H D F M F V K R M D P D C P H S D R I Y Y L A H S G E N T 229
 901 GAAAGGAGACTCACTTATGTCACAAATCAGCTAGCCAACTAGGAGAGATGCCAGATCAGCTGAGTCCCTACCTTTGTTCTGCAAGAGAAATTTGATA 1000
 230 E R R L T Y V H N E L A N H E E D A R S A G V A T F V L Q E E F D R 263
 1001 GATATTCCTGCTTATGCTGCTGCTCCAAAGCTGAAACAACTCCCACTGGTGGTAAATTTCTTGAATTTCTATGAAGAAATGATGAATCTGAGGTGA 1100
 263 Y S G Y W H C P K A E T T F S G G K I L R I L Y E E N D E S E V E 296
 1101 AATTATTCATGTTACATCCCTATGTTGAAACAGGAGGCGAGATTCATCCCTTATCTTAAACAGGTACAGCAAACTCTAAAGTCACTTTTAAGATG 1200
 296 I I H V T S P K L E T R R A D S F R Y P K T G T A M P K V T F K H 329
 1201 TCAGAAATATGATGATGCTGAGGAGGATCATAGATGTCATAGGAGGAGTCACTTCAACTTTTACAGATTCTATTTGAAGAGTTCGAATATATG 1300
 330 S E I M I D A E G I I D V I D K E L I Q F P E I L F E G V E Y I A 363
 1301 CCAGAGCTGGATGGATCTCTGAGGAGAAATATGCTTGGTCCATCTCTAGATAGCTCCAGACTGCGCTACAGATAGTGTGATCTCACTGAATATT 1400
 363 R A G W T P E G E K Y A W S I L L D R S O T R L O I V L I S P E L F 396
 1401 TATCCAGTAGAAGATGATGTTATGGAAGGCGAGAGCTCATTGAGTCAGTCCCTGATCTCTGAGGCGCACTAATATCTATGAAGAAACAGACATC 1500
 396 I P V E D D V M E R Q L I E S V F D S V T F L I I Y E E T T D I 429
 1501 TGGATAAATATCCATGACATCTTTATGTTTTCCTCCAAAGTCCAGAGGAGAAATGAGTTTATTTTGGCTCTGAATGCAAAAGAGTTCCTGCAAT 1600
 430 W I N I H D I F H V F P Q S H E E E I E F I F A S E C K T G F R H L 463
 1601 TATACAAATATCATCTATTTTAAAGGAGGAGAAATATAGGAGTCCAGTGGTGGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCT 1700
 463 Y K I T S I L K E S K Y K R S S G G L P A P S D F K C P I K E E I 496
 1701 AGCAATACCAGTGGTGAATGGAGGAGTCTTGGCGCGGATGGATCTAATATCCAGGTTGATGAAGTCAGAGGCTGGTATATTTTGAAGGACCAAGAGC 1800
 496 A I T S G E W E V L G R H G S N I O V D E V R R L V Y F E G T K 529
 1801 TCCCTTTAGAGCATCACTCTGAT 1900
 530 S P L E H H L Y V V S Y V N P G E V T R L T D R G Y S H S C C I S O 563
 1901 AGCACTGCTGCTCTTTTATAGTAAATATAGTAAACAGGAGATCCAGACTGCTGCTGCTTACAGCTATCAAGTCTGAGATGAGGAGGAGTCCGAA 2000
 563 H C D F F I S K Y S M Q R N P H C V S L Y K L S S P E D D P T C K 596
 2001 AACAAAGGAATTTTGGGCGGAGTTCGAT 2100
 596 T K E F W A T I L D S A G F L P D Y T P P E I F S P E S T T C F T 629
 2101 TGTATGGATGCTCTACAGGCTCATGATCTACAGGCTGGAAGAAATATCTACTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCT 2200
 630 L Y G H L Y K P M D L Q P G E K Y P T V L F I Y G G P Q V L V N H 663
 2201 ATCGGTTTAAAGGAGTCAAGTATTTCCCTTGAATACCTGAGCTCTCTGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGAT 2300
 663 R F K G V K Y P R L N T L A S L G Y V V V V I D N R G S C H R G L 696
 2301 TAAATTTGAAGGCGCTTTTAAATATAAATGAGTCAATAGAAATGAGGAGTCAAGTGGAGGAGTCCAAATATCTAGCTCTGATGATGATGATGATGAT 2400
 696 K F E G A F K Y K H G Q I E I D D O V E G L Q Y L A S R Y D F I D 729
 2401 TTAGATGCT 2500
 730 L D R V C I N G Y L S L H A L M Q R S D I P R V A I A G A 763
 2501 CCCCAGTCACTCTGATGATCTTCTATGATACAGGATACAGGAGGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGAT 2600
 763 P V T L W I F Y D T G Y T E R Y K H G P D O N E Q G Y L G S V A 796
 2601 CATGCAAGCAGAAAGTCCCTCTCAACCAATGCTTACTGCTCTTACATGCTTCTGATGATGATGATGATGATGATGATGATGATGATGATGATGAT 2700
 796 M Q A E K F P S E P M R L L L L H G F L B E N V H P A T S I L L 829
 2701 AGTTTTTAGTCAAGGCTGCAAGGAT 2800
 830 S F L V R A G K P Y D L Q I Y P O E R B S I R V P E S G E N Y E L H 863
 2801 ATCTTTGCACTCACTCTCAAGAAAGCTTGGATCAGTATGCTGCTCTAAAGGATGATGATGATGATGATGATGATGATGATGATGATGATGATGAT 2900
 863 L L H Y L Q E N L G S R I A L K V I 896
 2901 TTTAACCAGATGAGGAGTTTAAATCAACAGAAACACAGAAATGATCAGATTTTATGATCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCT 3000
 3001 CCATGAGGCGCTACCGTTTGTGAT 3100
 3101 AGAATTACTAAAAA 3127

FIGURE 2

3/9

bdpp8	HNHFYQFLYLQVSPFIDFSV	MAAANE	ELQCLV	I	FET	ADCE	WIE	25
cedpp8	HNHFYQFLYLQVSPFIDFSV	MAAANE	ELQCLV	I	FET	ADCE	WIE	50
bdpp4	HNHFYQFLYLQVSPFIDFSV	MAAANE	ELQCLV	I	FET	ADCE	WIE	0
bdpp8	SQDRFELPFFVYERY	SQLEKE	LADTRKY	UCTHMAKAP	ND			66
cedpp8	SQDRFELPFFVYERY	SQLEKE	LADTRKY	UCTHMAKAP	ND			100
bdpp4	SQDRFELPFFVYERY	SQLEKE	LADTRKY	UCTHMAKAP	ND			22
bdpp8	FLFLKNDTQCPH	SDRIYTLAK	RSGENRZNTLFYS	QI	PKTI	NR	AVTGL	114
cedpp8	FLFLKNDTQCPH	SDRIYTLAK	RSGENRZNTLFYS	QI	PKTI	NR	AVTGL	146
bdpp4	FLFLKNDTQCPH	SDRIYTLAK	RSGENRZNTLFYS	QI	PKTI	NR	AVTGL	51
bdpp8	SHPLLDLQATLD	YTC	MY	SPZ	LRL	GR	KXCTV	151
cedpp8	SHPLLDLQATLD	YTC	MY	SPZ	LRL	GR	KXCTV	192
bdpp4	SHPLLDLQATLD	YTC	MY	SPZ	LRL	GR	KXCTV	101
bdpp8	SYDYHQCSTFLFOAG	SC	YV	YD	GC	FP	GF	196
cedpp8	SYDYHQCSTFLFOAG	SC	YV	YD	GC	FP	GF	242
bdpp4	SYDYHQCSTFLFOAG	SC	YV	YD	GC	FP	GF	151
bdpp8	IRNDPRLCPADPD	WIAFI	SNDIN	SK	VI	TR	229	
cedpp8	IRNDPRLCPADPD	WIAFI	SNDIN	SK	VI	TR	288	
bdpp4	IRNDPRLCPADPD	WIAFI	SNDIN	SK	VI	TR	188	
bdpp8	ERRLTYVSHZLANH	ED	AKSA	CVAT	FV	LQEE	DRYSCY	278
cedpp8	ERRLTYVSHZLANH	ED	AKSA	CVAT	FV	LQEE	DRYSCY	317
bdpp4	ERRLTYVSHZLANH	ED	AKSA	CVAT	FV	LQEE	DRYSCY	222
bdpp8	CGKILLLYEZFV	SPZ	LY	YTS	PH	LT	ARADSP	325
cedpp8	CGKILLLYEZFV	SPZ	LY	YTS	PH	LT	ARADSP	361
bdpp4	CGKILLLYEZFV	SPZ	LY	YTS	PH	LT	ARADSP	266
bdpp8	THNHSZINIDAE	CRXZ	IV	ID	REL	LD	QF	372
cedpp8	THNHSZINIDAE	CRXZ	IV	ID	REL	LD	QF	403
bdpp4	THNHSZINIDAE	CRXZ	IV	ID	REL	LD	QF	297
bdpp8	YARSTLDRSDQ	AL	LY	LY	LY	LY	LY	400
cedpp8	YARSTLDRSDQ	AL	LY	LY	LY	LY	LY	453
bdpp4	YARSTLDRSDQ	AL	LY	LY	LY	LY	LY	330
SPBMCDY3-3-10								
bdpp8	DDVNEQRXZSV	DSV	TL	LI	YE	ET	DI	447
cedpp8	DDVNEQRXZSV	DSV	TL	LI	YE	ET	DI	501
bdpp4	DDVNEQRXZSV	DSV	TL	LI	YE	ET	DI	369
bdpp8	YEFIFA	SE	CR	IT	GL	LY	HT	495
cedpp8	YEFIFA	SE	CR	IT	GL	LY	HT	536
bdpp4	YEFIFA	SE	CR	IT	GL	LY	HT	396
bdpp8	IAISGZNPV	LG	EC	SH	VO	DE	VAL	543
cedpp8	IAISGZNPV	LG	EC	SH	VO	DE	VAL	581
bdpp4	IAISGZNPV	LG	EC	SH	VO	DE	VAL	435
bdpp8	ICZVTRLTDR	CYS	ESC	CL	SA	CD	FF	585
cedpp8	ICZVTRLTDR	CYS	ESC	CL	SA	CD	FF	629
bdpp4	ICZVTRLTDR	CYS	ESC	CL	SA	CD	FF	481
bdpp8	LSSELDPTCKT	FE	RA	IT	LD	SA	CP	591
cedpp8	LSSELDPTCKT	FE	RA	IT	LD	SA	CP	628
bdpp4	LSSELDPTCKT	FE	RA	IT	LD	SA	CP	576
bdpp8	CHLH	FE	RA	IT	LD	SA	CP	601
cedpp8	CHLH	FE	RA	IT	LD	SA	CP	625
bdpp4	CHLH	FE	RA	IT	LD	SA	CP	572
bdpp8	YVY	FE	RA	IT	LD	SA	CP	629
cedpp8	YVY	FE	RA	IT	LD	SA	CP	624
bdpp4	YVY	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH	FE	RA	IT	LD	SA	CP	624
bdpp4	HLH	FE	RA	IT	LD	SA	CP	520
bdpp8	HLH	FE	RA	IT	LD	SA	CP	629
cedpp8	HLH							




FIGURE 4





FIGURE 5

6/9

**FIGURE 6**

7/9




FIGURE 7




FIGURE 8

FIGURE 9A





FIGURE 9B



1.ST25.txt
SEQUENCE LISTING

<110> THE UNIVERSITY OF SYDNEY

<120> DIPEPTIDYL PEPTIDASE

<130> P37354

<150> AU PQ2762

<151> 1999-09-10

<150> AU PQ5709

<151> 2000-02-18

<160> 8

<170> PatentIn version 3.0

<210> 1

<211> 882

<212> PRT

<213> Homo sapiens

<400> 1

Met	Ala	Ala	Ala	Met	Glu	Thr	Glu	Gln	Leu	Gly	Val	Glu	Ile	Phe	Glu	1	5	10	15
Thr	Ala	Asp	Cys	Glu	Glu	Asn	Ile	Glu	Ser	Gln	Asp	Arg	Pro	Lys	Leu	20	25	30	
Glu	Pro	Phe	Tyr	Val	Glu	Arg	Tyr	Ser	Trp	Ser	Gln	Leu	Lys	Lys	Leu	35	40	45	
Leu	Ala	Asp	Thr	Arg	Lys	Tyr	His	Gly	Tyr	Met	Met	Ala	Lys	Ala	Pro	50	55	60	
His	Asp	Phe	Met	Phe	Val	Lys	Arg	Asn	Asp	Pro	Asp	Gly	Pro	His	Ser	65	70	75	80
Asp	Arg	Ile	Tyr	Tyr	Leu	Ala	Met	Ser	Gly	Glu	Asn	Arg	Glu	Asn	Thr	85	90	95	
Leu	Phe	Tyr	Ser	Glu	Ile	Pro	Lys	Thr	Ile	Asn	Arg	Ala	Ala	Val	Leu	100	105	110	
Met	Leu	Ser	Trp	Lys	Pro	Leu	Leu	Asp	Leu	Phe	Gln	Ala	Thr	Leu	Asp	115	120	125	
Tyr	Gly	Met	Tyr	Ser	Arg	Glu	Glu	Glu	Leu	Leu	Arg	Glu	Arg	Lys	Arg	130	135	140	
Ile	Gly	Thr	Val	Gly	Ile	Ala	Ser	Tyr	Asp	Tyr	His	Gln	Gly	Ser	Gly	145	150	155	160

1.ST25.txt

Thr Phe Leu Phe Gln Ala Gly Ser Gly Ile Tyr His Val Lys Asp Gly
 165 170 175
 Gly Pro Gln Gly Phe Thr Gln Gln Pro Leu Arg Pro Asn Leu Val Glu
 180 185 190
 Thr Ser Cys Pro Asn Ile Arg Met Asp Pro Lys Leu Cys Pro Ala Asp
 195 200 205
 Pro Asp Trp Ile Ala Phe Ile His Ser Asn Asp Ile Trp Ile Ser Asn
 210 215 220
 Ile Val Thr Arg Glu Glu Arg Arg Leu Thr Tyr Val His Asn Glu Leu
 225 230 235 240
 Ala Asn Met Glu Glu Asp Ala Arg Ser Ala Gly Val Ala Thr Phe Val
 245 250 255
 Leu Gln Glu Glu Phe Asp Arg Tyr Ser Gly Tyr Trp Trp Cys Pro Lys
 260 265 270
 Ala Glu Thr Thr Pro Ser Gly Gly Lys Ile Leu Arg Ile Leu Tyr Glu
 275 280 285
 Glu Asn Asp Glu Ser Glu Val Glu Ile Ile His Val Thr Ser Pro Met
 290 295 300
 Leu Glu Thr Arg Arg Ala Asp Ser Phe Arg Tyr Pro Lys Thr Gly Thr
 305 310 315 320
 Ala Asn Pro Lys Val Thr Phe Lys Met Ser Glu Ile Met Ile Asp Ala
 325 330 335
 Glu Gly Arg Ile Ile Asp Val Ile Asp Lys Glu Leu Ile Gln Pro Phe
 340 345 350
 Glu Ile Leu Phe Glu Gly Val Glu Tyr Ile Ala Arg Ala Gly Trp Thr
 355 360 365
 Pro Glu Gly Lys Tyr Ala Trp Ser Ile Leu Leu Asp Arg Ser Gln Thr
 370 375 380
 Arg Leu Gln Ile Val Leu Ile Ser Pro Glu Leu Phe Ile Pro Val Glu
 385 390 395 400
 Asp Asp Val Met Glu Arg Gln Arg Leu Ile Glu Ser Val Pro Asp Ser
 405 410 415
 Val Thr Pro Leu Ile Ile Tyr Glu Glu Thr Thr Asp Ile Trp Ile Asn
 420 425 430
 Ile His Asp Ile Phe His Val Phe Pro Gln Ser His Glu Glu Glu Ile
 435 440 445
 Glu Phe Ile Phe Ala Ser Glu Cys Lys Thr Gly Phe Arg His Leu Tyr

1.ST25.txt

450					455					460					
Lys 465	Ile	Thr	Ser	Ile	Leu 470	Lys	Glu	Ser	Lys	Tyr 475	Lys	Arg	Ser	Ser	Gly 480
Gly	Leu	Pro	Ala	Pro 485	Ser	Asp	Phe	Lys	Cys 490	Pro	Ile	Lys	Glu	Glu 495	Ile
Ala	Ile	Thr	Ser 500	Gly	Glu	Trp	Glu	Val 505	Leu	Gly	Arg	His	Gly 510	Ser	Asn
Ile	Gln	Val 515	Asp	Glu	Val	Arg	Arg 520	Leu	Val	Tyr	Phe	Glu 525	Gly	Thr	Lys
Asp 530	Ser	Pro	Leu	Glu	His	His 535	Leu	Tyr	Val	Val	Ser 540	Tyr	Val	Asn	Pro
Gly 545	Glu	Val	Thr	Arg	Leu 550	Thr	Asp	Arg	Gly	Tyr 555	Ser	His	Ser	Cys	Cys 560
Ile	Ser	Gln	His	Cys 565	Asp	Phe	Phe	Ile	Ser 570	Lys	Tyr	Ser	Asn	Gln 575	Lys
Asn	Pro	His	Cys 580	Val	Ser	Leu	Tyr	Lys 585	Leu	Ser	Ser	Pro	Glu 590	Asp	Asp
Pro	Thr	Cys 595	Lys	Thr	Lys	Glu	Phe 600	Trp	Ala	Thr	Ile	Leu 605	Asp	Ser	Ala
Gly 610	Pro	Leu	Pro	Asp	Tyr	Thr 615	Pro	Pro	Glu	Ile	Phe 620	Ser	Phe	Glu	Ser
Thr 625	Thr	Gly	Phe	Thr	Leu 630	Tyr	Gly	Met	Leu	Tyr 635	Lys	Pro	His	Asp	Leu 640
Gln	Pro	Gly	Lys	Lys 645	Tyr	Pro	Thr	Val	Leu 650	Phe	Ile	Tyr	Gly	Gly 655	Pro
Gln	Val	Gln 660	Leu	Val	Asn	Asn	Arg	Phe 665	Lys	Gly	Val	Lys	Tyr 670	Phe	Arg
Leu	Asn 675	Thr	Leu	Ala	Ser	Leu	Gly 680	Tyr	Val	Val	Val	Val 685	Ile	Asp	Asn
Arg 690	Gly	Ser	Cys	His	Arg	Gly 695	Leu	Lys	Phe	Glu	Gly 700	Ala	Phe	Lys	Tyr
Lys 705	Met	Gly	Gln	Ile	Glu 710	Ile	Asp	Asp	Gln	Val 715	Glu	Gly	Leu	Gln 720	Tyr
Leu	Ala	Ser	Arg	Tyr 725	Asp	Phe	Ile	Asp	Leu 730	Asp	Arg	Val	Gly	Ile 735	His
Gly	Trp	Ser	Tyr 740	Gly	Gly	Tyr	Leu	Ser	Leu 745	Met	Ala	Leu	Met 750	Gln	Arg

1.ST25.txt

Ser Asp Ile Phe Arg Val Ala Ile Ala Gly Ala Pro Val Thr Leu Trp
755 760 765

Ile Phe Tyr Asp Thr Gly Tyr Thr Glu Arg Tyr Met Gly His Pro Asp
770 775 780

Gln Asn Glu Gln Gly Tyr Tyr Leu Gly Ser Val Ala Met Gln Ala Glu
785 790 795 800

Lys Phe Pro Ser Glu Pro Asn Arg Leu Leu Leu Leu His Gly Phe Leu
805 810 815

Asp Glu Asn Val His Phe Ala His Thr Ser Ile Leu Leu Ser Phe Leu
820 825 830

Val Arg Ala Gly Lys Pro Tyr Asp Leu Gln Ile Tyr Pro Gln Glu Arg
835 840 845

His Ser Ile Arg Val Pro Glu Ser Gly Glu His Tyr Glu Leu His Leu
850 855 860

Leu His Tyr Leu Gln Glu Asn Leu Gly Ser Arg Ile Ala Ala Leu Lys
865 870 875 880

Val Ile

<210> 2

<211> 3120

<212> DNA

<213> Homo sapiens

<400> 2

aagtgtctaaa gcctccgagg ccaaggccgc tgctactgcc gccgctgctt cttagtgccg
60

cgttcgccgc ctgggttggtc accggcgccg ccgcccaggga agccactgca accaggaccg
120

gagtggaggc ggcgagcat gaagcggcgc aggcccgctc catagcgcac gtcgggacgg
180

tccggggcggg gccgggggga aggaaaatgc aacatggcag cagcaatgga aacagaacag
240

ctgggtgttg agatatttga aactgcggac tgtgaggaga atattgaatc acaggatcgg
300

cctaaattgg agccttttta tggtgagcgg tattcctgga gtcagcttaa aaagctgctt
360

gccgatacca gaaaatatca tggctacatg atggctaagg caccacatga tttcatgttt
420

1.ST25.txt

gtgaagagga atgatccaga tggacctcat tcagacagaa tctattacct tgccatgtct
480

ggtgagaaca gagaaaatac actgttttat tctgaaattc ccaaaactat caatagagca
540

gcagtcttaa tgctctcttg gaagcctctt ttggatcttt ttcaggcaac actggactat
600

ggaatgtatt ctcgagaaga agaactatta agagaaagaa aacgcattgg aacagtcgga
660

attgcttctt acgattatca ccaaggaagt ggaacatttc tgtttcaagc cggtagtgga
720

atttatcacg taaaagatgg agggccacaa ggatttacgc aacaaccttt aaggcccaat
780

ctagtggaaa ctagtgtgcc caacatacgg atggatccaa aattatgccc cgctgatcca
840

gactggattg cttttataca tagcaacgat atttggatat ctaacatcgt aaccagagaa
900

gaaaggagac tcacttatgt gcacaatgag ctagccaaca tggaagaaga tgccagatca
960

gctggagtcg ctacctttgt tctccaagaa gaatttgata gatattctgg ctattggtgg
1020

tgtccaaaag ctgaaacaac tcccagtggg ggtaaaattc ttagaattct atatgaagaa
1080

aatgatgaat ctgagggtgga aattattcat gttacatccc ctatgttgga aacaaggagg
1140

gcagattcat tccgttatcc taaaacaggt acagcaaacc ctaaagtcac ttttaagatg
1200

tcagaaataa tgattgatgc tgaaggaagg atcatagatg tcatagataa ggaactaatt
1260

caaccttttg agattctatt tgaaggagtt gaatatattg ccagagctgg atggactcct
1320

gagggaaaat atgcttggtc catcctacta gatcgctccc agactcgctt acagatagtg
1380

ttgatctcac ctgaattatt tatcccagta gaagatgatg ttatggaaag gcagagactc
1440

attgagtcag tgcctgattc tgtgacgcca ctaattatct atgaagaaac aacagacatc
1500

tggataaata tccatgacat ctttcatggt ttccccaaa gtcacgaaga ggaaattgag

1.ST25.txt

1560

tttatttttg cctctgaatg caaaacaggt ttccgtcatt tatacaaaat tacatctatt
1620

ttaaaggaaa gcaaataata acgatccagt ggtgggctgc ctgctccaag tgatttcaag
1680

tgctctatca aagaggagat agcaattacc agtgggtgaat ggggaagttct tggccggcat
1740

ggatctaata tccaagttga tgaagtcaga aggctggat attttgaagg caccaaagac
1800

tccccttttag agcatcacct gtacgtagtc agttacgtaa atcctggaga ggtgacaagg
1860

ctgactgacc gtggctactc acattcttgc tgcatacgtc agcactgtga cttctttata
1920

agtaagtata gtaaccagaa gaatccacac tgtgtgtccc tttacaagct atcaagtcct
1980

gaagatgacc caacttgcaa aacaaaggaa ttttgggcca ccattttgga ttcagcaggt
2040

cctcttcttg actatactcc tccagaaatt ttctcttttg aaagtactac tggatttaca
2100

ttgtatggga tgctctacaa gcctcatgat ctacagcctg gaaagaaata tcctactgtg
2160

ctgttcatat atgggtggtec tcaggtgcag ttgggtgaata atcggtttta aggagtcaag
2220

tatttccgct tgaataccct agcctctcta gggtatgtgg ttgtagtgat agacaacagg
2280

ggatcctgtc accgagggct taaatttgaa ggcgccttta aatataaaat gggtaaata
2340

gaaattgacg atcaggtgga aggactccaa tatctagctt ctcgatatga tttcattgac
2400

ttagatcgtg tgggcatcca cggctggtec tatggaggat acctctccct gatggcatta
2460

atgcagaggt cagatatctt cagggttgct attgctgggg cccaggtcac tctgtggatc
2520

ttctatgata caggatacac ggaacgttat atgggtcacc ctgaccagaa tgaacagggc
2580

tattacttag gatctgtggc catgcaagca gaaaagttcc cctctgaacc aaatcgttta
2640

1.ST25.txt

ctgctcttac atggtttctt ggatgagaat gtccattttg cacataccag tatattactg
2700

agtttttttag tgagggttg aaagccatat gatttacaga tctatcctca ggagagacac
2760

agcataagag ttcttgaatc gggagaacat tatgaactgc atcttttgca ctaccttcaa
2820

gaaaaccttg gatcacgtat tgctgctcta aaagtgatat aattttgacc tgtgtagaac
2880

tctctggtat acactggcta tttaaccaaa tgaggagggtt taatcaacag aaaacacaga
2940

attgatcatc acattttgat acctgccatg taacatctac tctgaaaat aaatgtggtg
3000

ccatgcaggg gtctacggtt tgtggtagta atctaatacc ttaacccac atgctcaaaa
3060

tcaaatgata catattcctg agagaccag caataccata agaattacta aaaaaaaaaa
3120

<210> 3
<211> 310
<212> PRT
<213> Homo sapiens

<400> 3

Phe	Glu	Gly	Thr	Lys	Asp	Ser	Pro	Leu	Glu	His	His	Leu	Tyr	Val	Val
1				5					10					15	
Ser	Tyr	Val	Asn	Pro	Gly	Glu	Val	Thr	Arg	Leu	Thr	Asp	Arg	Gly	Tyr
			20					25					30		
Ser	His	Ser	Cys	Cys	Ile	Ser	Gln	His	Cys	Asp	Phe	Phe	Ile	Ser	Lys
			35				40					45			
Tyr	Ser	Asn	Gln	Lys	Asn	Pro	His	Cys	Val	Ser	Leu	Tyr	Lys	Leu	Ser
	50					55					60				
Ser	Pro	Glu	Asp	Asp	Pro	Thr	Cys	Lys	Thr	Lys	Glu	Phe	Trp	Ala	Thr
65					70				75					80	
Ile	Leu	Asp	Ser	Ala	Gly	Pro	Leu	Pro	Asp	Tyr	Thr	Pro	Pro	Glu	Ile
				85					90					95	
Phe	Ser	Phe	Glu	Ser	Thr	Thr	Gly	Phe	Thr	Leu	Tyr	Gly	Met	Leu	Tyr
			100					105					110		
Lys	Pro	His	Asp	Leu	Gln	Pro	Gly	Lys	Lys	Tyr	Pro	Thr	Val	Leu	Phe

1.ST25.txt

115	120	125
Ile Tyr Gly Gly Pro Gln Gly	Gln Ile Glu Ile Asp Asp Gln Val Glu	
130	135	140
Gly Leu Gln Tyr Leu Ala Ser Arg Tyr Asp Phe Ile Asp Leu Asp Arg		
145	150	155
Val Gly Ile His Gly Trp Ser Tyr Gly Gly Tyr Leu Ser Leu Met Ala		
165	170	175
Leu Met Gln Arg Ser Asp Ile Phe Arg Val Ala Ile Ala Gly Ala Pro		
180	185	190
Val Thr Leu Trp Ile Phe Tyr Asp Thr Gly Tyr Thr Glu Arg Tyr Met		
195	200	205
Gly His Pro Asp Gln Asn Glu Gln Gly Tyr Tyr Leu Gly Ser Val Ala		
210	215	220
Met Gln Ala Glu Lys Phe Pro Ser Glu Pro Asn Arg Leu Leu Leu Leu		
225	230	235
His Gly Phe Leu Asp Glu Asn Val His Phe Ala His Thr Ser Ile Leu		
245	250	255
Leu Ser Phe Leu Val Arg Ala Gly Lys Pro Tyr Asp Leu Gln Ile Tyr		
260	265	270
Pro Gln Glu Arg His Ser Ile Arg Val Pro Glu Ser Gly Glu His Tyr		
275	280	285
Glu Leu His Leu Leu His Tyr Leu Gln Glu Asn Leu Gly Ser Arg Ile		
290	295	300
Ala Ala Leu Lys Val Ile		
305	310	

<210> 4

<211> 1197

<212> DNA

<213> Homo sapiens

<400> 4

attttgaagg caccaaagac tccccttttag agcatcacct gtacgtagtc agttacgtaa
60

atcctggaga ggtgacaagg ctgactgacc gtggctactc acattcttgc tgcacagtc
120

agcactgtga cttctttata agtaagtata gtaaccagaa gaatccacac tgtgtgtccc
180

tttacaagct atcaagtcct gaagatgacc caacttgcaa aacaaaggaa ttttggggcca
240

1.ST25.txt

ccattttgga ttcagcaggt cctcttcctg actatactcc tccagaaatt ttctcttttg
300
aaagtactac tggatttaca ttgtatggga tgctctacaa gcctcatgat ctacagcctg
360
gaaagaaata tctactgtg ctgttcatat atgggtggcc tcagggtcaa atagaaattg
420
acgatcaggt ggaaggactc caatatctag cttctcgata tgatttcatt gacttagatc
480
gtgtgggcat ccacggctgg tcctatggag gatacctctc cctgatggca ttaatgcaga
540
ggtcagatat cttcagggtt gctattgctg gggccccagt cactctgtgg atcttctatg
600
atacaggata cacggaacgt tatatgggtc accctgacca gaatgaacag ggctattact
660
taggatctgt ggccatgcaa gcagaaaagt tcccctctga accaaatcgt ttactgctct
720
tacatggttt cctggatgag aatgtccatt ttgcacatac cagtatatta ctgagttttt
780
tagtgagggc tggaaagcca tatgatttac agatctatcc tcaggagaga cacagcataa
840
gagttcctga atcgggagaa cattatgaac tgcactcttt gcactacctt caagaaaacc
900
ttggatcacg tattgctgct ctaaaagtga tataattttg acctgtgtag aactctctgg
960
tatacactgg ctatttaacc aaatgaggag gtttaatcaa cagaaaacac agaattgatc
1020
atcacatttt gatacctgcc atgtaacatc tactcctgaa aataaatgtg gtgccatgca
1080
ggggtctacg gtttgtggta gtaatctaata accttaaccc cacatgctca aaatcaaattg
1140
atacatattc ctgagagacc cagcaatacc ataagaatta ctaaaaaaaaa aaaaaaa
1197

<210> 5
<211> 465
<212> PRT
<213> Homo sapiens

1..ST25.txt

<400> 5

```

Thr Gly Thr Ala Asn Pro Lys Val Thr Phe Lys Met Ser Glu Ile Met
1          5          10          15
Ile Asp Ala Glu Gly Arg Ile Ile Asp Val Ile Asp Lys Glu Leu Ile
20          25          30
Gln Pro Phe Glu Ile Leu Phe Glu Gly Val Glu Tyr Ile Ala Arg Ala
35          40          45
Gly Trp Thr Pro Glu Gly Lys Tyr Ala Trp Ser Ile Leu Leu Asp Arg
50          55          60
Ser Gln Thr Arg Leu Gln Ile Val Leu Ile Ser Pro Glu Leu Phe Ile
65          70          75          80
Pro Val Glu Asp Asp Val Met Glu Arg Gln Arg Leu Ile Glu Ser Val
85          90          95
Pro Asp Ser Val Thr Pro Leu Ile Ile Tyr Glu Glu Thr Thr Asp Ile
100         105         110
Trp Ile Asn Ile His Asp Ile Phe His Val Phe Pro Gln Ser His Glu
115         120         125
Glu Glu Ile Glu Phe Ile Phe Ala Ser Glu Cys Lys Thr Gly Phe Arg
130         135         140
His Leu Tyr Lys Ile Thr Ser Ile Leu Lys Glu Ser Lys Tyr Lys Arg
145         150         155         160
Ser Ser Gly Gly Leu Pro Ala Pro Ser Asp Phe Lys Cys Pro Ile Lys
165         170         175
Glu Glu Ile Ala Ile Thr Ser Gly Glu Trp Glu Val Leu Gly Arg His
180         185         190
Gly Ser Asn Ile Gln Val Asp Glu Val Arg Arg Leu Val Tyr Phe Glu
195         200         205
Gly Thr Lys Asp Ser Pro Leu Glu His His Leu Tyr Val Val Ser Tyr
210         215         220
Val Asn Pro Gly Glu Val Thr Arg Leu Thr Asp Arg Gly Tyr Ser His
225         230         235         240
Ser Cys Cys Ile Ser Gln His Cys Asp Phe Phe Ile Ser Lys Tyr Ser
245         250         255
Asn Gln Lys Asn Pro His Cys Val Ser Leu Tyr Lys Leu Ser Ser Pro
260         265         270
Glu Asp Asp Pro Thr Cys Lys Thr Lys Glu Phe Trp Ala Thr Ile Leu
275         280         285

```

1.ST25.txt

Asp Ser Ala Gly Pro Leu Pro Asp Tyr Thr Pro Pro Glu Ile Phe Ser
 290 295 300
 Phe Glu Ser Thr Thr Gly Phe Thr Leu Tyr Gly Met Leu Tyr Lys Pro
 305 310 315 320
 His Asp Leu Gln Pro Gly Lys Lys Tyr Pro Thr Val Leu Phe Ile Tyr
 325 330 335
 Gly Gly Pro Gln Val Ala Ile Ala Gly Ala Pro Val Thr Leu Trp Ile
 340 345 350
 Phe Tyr Asp Thr Gly Tyr Thr Glu Arg Tyr Met Gly His Pro Asp Gln
 355 360 365
 Asn Glu Gln Gly Tyr Tyr Leu Gly Ser Val Ala Met Gln Ala Glu Lys
 370 375 380
 Phe Pro Ser Glu Pro Asn Arg Leu Leu Leu Leu His Gly Phe Leu Asp
 385 390 395 400
 Glu Asn Val His Phe Ala His Thr Ser Ile Leu Leu Ser Phe Leu Val
 405 410 415
 Arg Ala Gly Lys Pro Tyr Asp Leu Gln Ile Tyr Pro Gln Glu Arg His
 420 425 430
 Ser Ile Arg Val Pro Glu Ser Gly Glu His Tyr Glu Leu His Leu Leu
 435 440 445
 His Tyr Leu Gln Glu Asn Leu Gly Ser Arg Ile Ala Ala Leu Lys Val
 450 455 460
 Ile
 465
 <210> 6
 <211> 1669
 <212> DNA
 <213> Homo sapiens
 <400> 6
 aacaggtaca gcaaataccta aagtcacttt taagatgtca gaaataatga ttgatgctga
 60
 aggaaggatc atagatgtca tagataagga actaattcaa ccttttgaga ttctatttga
 120
 aggagttgaa tatattgccca gagctggatg gactcctgag ggaaaatatg cttgggtccat
 180
 cctactagat cgctcccaga ctcgcctaca gatagtgttg atctcacctg aattatttat
 240

1.ST25.txt

cccagtagaa gatgatgtta tggaaaggca gagactcatt gagtcagtgc ctgattctgt
300

gacgccacta attatctatg aagaacaac agacatctgg ataaatatcc atgacatctt
360

tcatgttttt ccccaaagtc acgaagagga aattgagttt atttttgcct ctgaatgcaa
420

aacaggtttc cgtcatttat acaaaattac atctatttta aaggaaagca aatataaacg
480

atccagtggg gggctgcctg ctccaagtga tttcaagtgt cctatcaaag aggagatagc
540

aattaccagt ggtgaatggg aagttcttgg ccggcatgga tctaatatcc aagttgatga
600

agtcagaagg ctggtatatt ttgaaggcac caaagactcc cctttagagc atcacctgta
660

cgtagtcagt tacgtaaata ctggagaggt gacaaggctg actgaccgtg gctactcaca
720

ttcttgctgc atcagtcagc actgtgactt ctttataagt aagtatagta accagaagaa
780

tccacactgt gtgtcccttt acaagctatc aagtcctgaa gatgacccaa cttgcaaaac
840

aaaggaatth tgggccacca ttttggattc agcaggctct cttcctgact atactcctcc
900

agaaatthtc tcttttgaaa gtactactgg atttacattg tatgggatgc tctacaagcc
960

tcatgatcta cagcctggaa agaaatatcc tactgtgctg ttcatatatg gtggctctca
1020

ggttgctatt gctggggccc cagtcactct gtggatcttc tatgatacag gatacacgga
1080

acgttatatg ggccaccctg accagaatga acagggtat tacttaggat ctgtggccat
1140

gcaagcagaa aagttccctt ctgaaccaa tcgtttactg ctcttacatg gtttccctgga
1200

tgagaatgtc cattttgcac ataccagtat attactgagt tttttagtga gggctggaaa
1260

gccatatgat ttacagatct atcctcagga gagacacagc ataagagttc ctgaatcggg
1320

agaacattat gaactgcata ttttgacta ctttcaagaa aaccttggat cacgtattgc

1.ST25.txt

1380

tgctctaaaa gtgatataat ttgacctgt gtagaactct ctggtataca ctggctattt
1440

aaccaaatga ggaggtttta tcaacagaaa acacagaatt gatcatcaca ttttgatacc
1500

tgccatgtaa catctactcc tgaaaataaaa tgtggtgccca tgcagggggtc tacgggttgt
1560

ggtagtaatc taatacctta accccacatg ctcaaaatca aatgatacat attcctgaga
1620

gaccagcaa taccataaga attactaaaa aaaaaaaaaa aaaaaaaaaa
1669

<210> 7

<211> 360

<212> PRT

<213> Homo sapiens

<400> 7

Glu Glu Asp Ala Arg Ser Ala Gly Val Ala Thr Phe Val Leu Gln Glu
1 5 10 15

Glu Phe Asp Arg Tyr Ser Gly Tyr Trp Trp Cys Pro Lys Ala Glu Thr
20 25 30

Thr Pro Ser Gly Gly Lys Ile Leu Arg Ile Leu Tyr Glu Glu Asn Asp
35 40 45

Glu Ser Glu Val Glu Ile Ile His Val Thr Ser Pro Met Leu Glu Thr
50 55 60

Arg Arg Ala Asp Ser Phe Arg Tyr Pro Lys Thr Gly Thr Ala Asn Pro
65 70 75 80

Lys Val Thr Phe Lys Met Ser Glu Ile Met Ile Asp Ala Glu Gly Arg
85 90 95

Ile Ile Val Asp Glu Val Arg Arg Leu Val Tyr Phe Glu Gly Thr Lys
100 105 110

Asp Ser Pro Leu Glu His His Leu Tyr Val Val Ser Tyr Val Asn Pro
115 120 125

Gly Glu Val Thr Arg Leu Thr Asp Arg Gly Tyr Ser His Ser Cys Cys
130 135 140

Ile Ser Gln His Cys Asp Phe Phe Ile Ser Lys Tyr Ser Asn Gln Lys
145 150 155 160

1.ST25.txt

Asn Pro His Cys Val Ser Leu Tyr Lys Leu Ser Ser Pro Glu Asp Asp
 165 170 175
 Pro Thr Cys Lys Thr Lys Glu Phe Trp Ala Thr Ile Leu Asp Ser Ala
 180 185 190
 Gly Pro Leu Pro Asp Tyr Thr Pro Pro Glu Ile Phe Ser Phe Glu Ser
 195 200 205
 Thr Thr Gly Phe Thr Leu Tyr Gly Met Leu Tyr Lys Pro His Asp Leu
 210 215 220
 Gln Pro Gly Lys Lys Tyr Pro Thr Val Leu Phe Ile Tyr Gly Gly Pro
 225 230 235 240
 Gln Val Gln Leu Val Asn Asn Arg Phe Lys Gly Val Lys Tyr Phe Arg
 245 250 255
 Leu Asn Thr Leu Ala Ser Leu Gly Tyr Val Val Val Val Ile Asp Asn
 260 265 270
 Arg Gly Ser Cys His Arg Gly Leu Lys Phe Glu Gly Ala Phe Lys Tyr
 275 280 285
 Lys Met Gly Gln Ile Glu Ile Asp Asp Gln Val Glu Gly Leu Gln Tyr
 290 295 300
 Leu Ala Ser Arg Tyr Asp Phe Ile Asp Leu Asp Arg Val Gly Ile His
 305 310 315 320
 Gly Trp Ser Tyr Gly Gly Tyr Leu Ser Leu Met Ala Leu Met Gln Arg
 325 330 335
 Ser Asp Ile Phe Arg Val Ala Ile Ala Gly Ala Pro Val Thr Leu Trp
 340 345 350
 Ile Phe Tyr Asp Thr Gly Tyr Thr
 355 360
 <210> 8
 <211> 1083
 <212> DNA
 <213> Homo sapiens
 <400> 8
 ggaagaagat gccagatcag ctggagtcgc tacctttggt ctccaagaag aatttgatag
 60
 atattctggc tattggtggt gtccaaaagc tgaaacaact cccagtgggtg gtaaaattct
 120
 tagaattcta tatgaagaaa atgatgaatc tgaggtggaa attattcatg ttacatcccc
 180
 tatgttgga acaaggagg cagattcatt ccgttatcct aaaacaggta cagcaaattcc

1.ST25.txt

240

taaagtcact ttttaagatgt cagaaataat gattgatgct gaaggaagga tcatagttga
300

tgaagtcaga aggctgggtat attttgaagg caccaaagac tccccttttag agcatcacct
360

gtacgtagtc agttacgtaa atcctggaga ggtgacaagg ctgactgacc gtggctactc
420

acattcttgc tgcacagtc agcactgtga cttctttata agtaagtata gtaaccagaa
480

gaatccacac tgtgtgtccc ttacaagct atcaagtcct gaagatgacc caacttgcaa
540

aacaaaggaa ttttgggcca ccattttgga ttcagcaggt cctcttctctg actatactcc
600

tccagaaatt ttctcttttg aaagtactac tggatttaca ttgtatggga tgctctacaa
660

gcctcatgat ctacagcctg gaaagaaata tcctactgtg ctgttcatat atgggtggctc
720

tcagggtgcag ttggtgaata atcggtttta aggagtcaag tatttccgct tgaataccct
780

agcctctcta gggttatgtgg ttgtagtgat agacaacagg ggatcctgtc accgagggct
840

taaatttgaa ggcgccttta aatataaaat ggggtcaaata gaaattgacg atcagggtgga
900

aggactccaa tatctagctt ctcgatatga ttccattgac ttagatcgtg tgggcatcca
960

cggctggctc tatggaggat acctctccct gatggcatta atgcagaggt cagatatctt
1020

cagggttgct attgctgggg cccdagtcac tctgtggatc ttctatgata caggatacac
1080

gga
1083

INTERNATIONAL SEARCH REPORT

International application No.
PCT/AU00/01085

A. CLASSIFICATION OF SUBJECT MATTER												
Int. Cl. ⁷ : C07K 14/47, C12N 9/64, 15/12, C12Q 1/37, A61K 38/48, A61P 19/02, 31/18, 37/00, 37/06, G01N 33/53.												
According to International Patent Classification (IPC) or to both national classification and IPC												
B. FIELDS SEARCHED												
Minimum documentation searched (classification system followed by classification symbols)												
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched												
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) Database: STN File Reg, CA and ANGIS. CA: subsequences of claim 1; Angis: SEQ ID NOS: 1 and 2 and homology search.												
C. DOCUMENTS CONSIDERED TO BE RELEVANT												
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.										
P,X	GenPept, Accession Number BAA 91059, submitted 15 February 2000, T. Kawakami et al, "NEDO human cDNA sequencing project."	1-3, 16-19										
A	FEBS Letters, vol 458, pages 278-284, 1999, C.A. Abbott et al, "Two highly conserved glutamic acid residues in the predicted beta propeller domain of dipeptidyl peptidase IV are required for its enzyme activity."	1-27										
A	European Journal of Biochemistry, vol 239, pages 248-258, 1996, E. Jacotot et al, "Dipeptidyl-peptidase IV-beta, a novel form of cell-surface-expressed protein with dipeptidyl-peptidase IV activity."	1-27										
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex												
<p>* Special categories of cited documents:</p> <table border="0"> <tr> <td>"A" document defining the general state of the art which is not considered to be of particular relevance</td> <td>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</td> </tr> <tr> <td>"E" earlier application or patent but published on or after the international filing date</td> <td>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</td> </tr> <tr> <td>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</td> <td>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</td> </tr> <tr> <td>"O" document referring to an oral disclosure, use, exhibition or other means</td> <td>"&" document member of the same patent family</td> </tr> <tr> <td>"P" document published prior to the international filing date but later than the priority date claimed</td> <td></td> </tr> </table>			"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention	"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone	"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art	"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family	"P" document published prior to the international filing date but later than the priority date claimed	
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention											
"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone											
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art											
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family											
"P" document published prior to the international filing date but later than the priority date claimed												
Date of the actual completion of the international search 16 October 2000		Date of mailing of the international search report 18 OCT 2000										
Name and mailing address of the ISA/AU AUSTRALIAN PATENT OFFICE PO BOX 200, WODEN ACT 2606, AUSTRALIA E-mail address: pct@ipaustalia.gov.au Facsimile No. (02) 6285 3929		Authorized officer <i>Frances Roden</i> FRANCES RODEN Telephone No: (02) 6283 2239										

INTERNATIONAL SEARCH REPORT

International application No.

PCT/AU00/01085

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,A	Immunological Reviews, vol 174, pages 172-191, 2000, G.W. McCaughan, "Molecular pathogenesis of liver disease: an approach to hepatic inflammation, cirrhosis and liver transplant tolerance."	1-27